

Group analysis of three dimensional Euler equations of gas dynamics

Mehdi Nadjafikhah*

Abstract

In this paper, the equations governing the unsteady flow of a perfect polytropic gas in three space dimensions are considered. The basic similarity reductions for this system are performed. Reduced equations and exact solutions associated with the symmetries are obtained. This results is used to give the invariance of system up to Galilean motions of space-times $\mathbb{R}^4$. Then, an optimal system of one-dimensional sub-algebras for symmetry algebra of this system is given.

Keywords: Euler equations, Lie group of transformations, Symmetry generators, Similarity solution, optimal system of Lie sub-algebras.

Mathematics Subject Classification: 35C05, 76M60.

1 Introduction

The equations governing the unsteady flow of a perfect polytropic gas in three space dimensions are

$$\begin{aligned} \frac{\partial u}{\partial t} + u \frac{\partial u}{\partial x} + v \frac{\partial u}{\partial y} + w \frac{\partial u}{\partial z} + \frac{1}{q} \frac{\partial p}{\partial x} &= 0, \\ \frac{\partial v}{\partial t} + u \frac{\partial v}{\partial x} + v \frac{\partial v}{\partial y} + w \frac{\partial v}{\partial z} + \frac{1}{q} \frac{\partial p}{\partial y} &= 0, \\ \frac{\partial w}{\partial t} + u \frac{\partial w}{\partial x} + v \frac{\partial w}{\partial y} + w \frac{\partial w}{\partial z} + \frac{1}{q} \frac{\partial p}{\partial z} &= 0, \\ \frac{\partial q}{\partial t} + q \left(\frac{\partial u}{\partial x} + \frac{\partial v}{\partial y} + \frac{\partial w}{\partial z} \right) + u \frac{\partial q}{\partial x} + v \frac{\partial q}{\partial y} + w \frac{\partial q}{\partial z} &= 0, \\ \frac{\partial p}{\partial t} + \gamma p \left(\frac{\partial u}{\partial x} + \frac{\partial v}{\partial y} + \frac{\partial w}{\partial z} \right) + u \frac{\partial p}{\partial x} + v \frac{\partial p}{\partial y} + w \frac{\partial p}{\partial z} &= 0. \end{aligned} \tag{1}$$

where t is the time and x, y, z are the space coordinates; $q(t, x, y, z)$ is the density, $p(t, x, y, z)$ the pressure, $u(t, x, y, z)$, $v(t, x, y, z)$ and $w(t, x, y, z)$

*Faculty of Mathematics, Iran University of Science and Technology, Narmak, Tehran, I.R.IRAN. e-mail: m_nadjafikhah@iust.ac.ir

the velocity components in the x , y and z directions, respectively, and $\gamma \in \mathbb{R}$ is the adiabatic index. These equations are called "*three dimensional Euler equations of gas dynamics*" [8] and section 6.3 of [9].

In this paper, we consider the equations governing the unsteady flow of a perfect polytropic gas in three space dimensions. The basic similarity reductions for this system are performed. Reduced equations and exact solutions associated with the symmetries are obtained. We find an optimal system of one-dimensional sub-algebras for symmetry algebra of this system.

In section 2 we find the full symmetry algebra of system (1). Chapter 3 is devoted to the group-invariant solutions to the system. The structure of full symmetry algebra illustrated in section. The last section deals with a optimal system of sub-algebras.

2 Lie symmetry of the system

The classical method for finding symmetry reductions of PDE is the Lie group method of infinitesimal transformations. To apply the classical method to (1), we consider the one-parameter Lie group of infinitesimal transformations in (t, x, y, z, u, v, p, q) given by

$$\begin{aligned}\tilde{t} &= t + s.\xi_1(t, x, y, z, u, v, p, q) + O(s^2), \\ \tilde{x} &= x + s.\xi_2(t, x, y, z, u, v, p, q) + O(s^2), \\ \tilde{y} &= y + s.\xi_3(t, x, y, z, u, v, p, q) + O(s^2), \\ \tilde{z} &= z + s.\xi_4(t, x, y, z, u, v, p, q) + O(s^2), \\ \tilde{u} &= u + s.\eta_1(t, x, y, z, u, v, p, q) + O(s^2), \\ \tilde{v} &= v + s.\eta_2(t, x, y, z, u, v, p, q) + O(s^2), \\ \tilde{w} &= w + s.\eta_3(t, x, y, z, u, v, p, q) + O(s^2), \\ \tilde{p} &= p + s.\eta_4(t, x, y, z, u, v, p, q) + O(s^2), \\ \tilde{q} &= q + s.\eta_5(t, x, y, z, u, v, p, q) + O(s^2).\end{aligned}\tag{2}$$

where s is the group parameter. One requires that this transformation leaves the sub-manifold S invariant. Let

$$X = \xi_1 \partial_t + \xi_2 \partial_x + \xi_3 \partial_y + \xi_4 \partial_z + \eta_1 \partial_u + \eta_2 \partial_v + \eta_3 \partial_w + \eta_4 \partial_p + \eta_5 \partial_q$$

be the corresponding infinitesimal transformation; where coefficients are real-valued C^∞ -functions of $t, x, y, z, u, v, w, p, q$. The infinitesimal criterion of invariance (1) will thus involve $t, x, y, z, u, v, w, p, q$ and the derivatives of u, v, w, p, q with respect to t, x, y, z , as well as $\xi_i, i = 1, 2, 3, 4$, and $\eta_j, j = 1, 2, 3, 4, 5$, and their partial derivatives. After eliminating any dependencies among the derivatives of the m 's caused by the system itself, we can then equate the coefficients of the remaining unconstrained partial derivatives of u to zero. This will result in a large number of elementary partial differential equations for the coefficient functions ξ_i, η_j of the infinitesimal generator, called the determining equations for the symmetry

group of the given system. In this case, we find the determining equations for the symmetry group of the system (1) to be the following:

$$\begin{aligned}
\xi_{1,t^2} &= \xi_{1,x} = \xi_{1,y} = \xi_{1,z} = \xi_{1,u} = \xi_{1,v} = \xi_{1,w} = \xi_{1,q} = \xi_{1,p} = 0, \\
\xi_{2,x} &= \xi_{2,u} = \xi_{2,v} = \xi_{2,w} = \xi_{2,q} = \xi_{2,p} = \xi_{2,t^2} = \xi_{2,ty} = \xi_{2,y^2} = 0, \\
\xi_{3,y} &= \xi_{3,u} = \xi_{3,v} = \xi_{3,w} = \xi_{3,q} = \xi_{3,p} = \xi_{3,t^2} = 0, \\
\xi_{4,z} &= \xi_{4,w} = \xi_{4,q} = \xi_{4,p} = 0, \\
\xi_{4,t^2} &= \xi_{4,tx} = \xi_{4,ty} = \xi_{4,x^2} = \xi_{4,xy} = \xi_{4,y^2} = 0, \\
\xi_{3,x} &= \xi_{2,y}, \quad \xi_{3,z} + \xi_{4,y} = 0, \quad \xi_{4,v} = \xi_{4,u}, \quad \xi_{2,z} + \xi_{4,x} = 0,
\end{aligned} \tag{3}$$

and

$$\begin{aligned}
\eta_1 &= -u\xi_{1,t} + \xi_{2,t} + v\xi_{2,y} - w\xi_{4,x}, \\
\eta_2 &= -w\xi_{4,y} + \xi_{3,t} - u\xi_{2,y} - v\xi_{1,t}, \\
\eta_3 &= u\xi_{4,x} + \xi_{4,t} + v\xi_{4,y} - w\xi_{1,t}, \\
\eta_4 &= q\eta_5/p + 2q\xi_{1,t}, \\
\eta_{5,t} &= \eta_{5,x} = \eta_{5,y} = \eta_{5,z} = \eta_{5,u} = \eta_{5,v} = \eta_{5,w} = \eta_{5,q} = 0, \\
\eta_5 &= p\eta_{5,p}.
\end{aligned} \tag{4}$$

First, equations (3) require that ξ_i s are just

$$\begin{aligned}
\xi_1 &= a_1t + a_2, \\
\xi_2 &= a_3t + a_4x + a_5y + a_6z + a_7, \\
\xi_3 &= a_8t - a_5x + a_4y + a_9z + a_{10}, \\
\xi_4 &= a_{11}t - a_6x - a_9y + a_4z + a_{12},
\end{aligned} \tag{5}$$

where $a_1, \dots, a_{12}$ are arbitrary constants. Then, equations (4) and (5) requires that

$$\begin{aligned}
\eta_1 &= (a_4 - a_1)u + a_5v - a_6w + a_3, \\
\eta_2 &= -a_5u + (a_4 - a_1)v + a_4w + a_8, \\
\eta_3 &= -a_6u - a_9v + (a_4 - a_1)w + a_{11}, \\
\eta_4 &= (2a_1 - 2a_4 + a_{13})q, \\
\eta_5 &= a_{13}p,
\end{aligned} \tag{6}$$

where a_{13} is an arbitrary constant. Therefore

Theorem 1. *The Lie algebra of infinitesimal symmetries of the system (1) is spanned by the 13 vector fields*

$$\begin{aligned}
X_1 &= \partial_x, \\
X_2 &= \partial_y, \\
X_3 &= \partial_z, \\
X_4 &= \partial_t, \\
X_5 &= t \partial_x + \partial_u, \\
X_6 &= t \partial_y + \partial_v, \\
X_7 &= t \partial_z + \partial_w, \\
X_8 &= t \partial_t - u \partial_u - v \partial_v - w \partial_w + 2q \partial_q, \\
X_9 &= x \partial_x + y \partial_y + z \partial_z + u \partial_u + v \partial_v + w \partial_w - 2r \partial_r, \\
X_{10} &= y \partial_x - x \partial_y + v \partial_u - u \partial_v, \\
X_{11} &= -z \partial_y + y \partial_z - w \partial_v + v \partial_w, \\
X_{12} &= -z \partial_x + x \partial_z - w \partial_u + u \partial_w, \\
X_{13} &= q \partial_q + p \partial_p,
\end{aligned}
\tag{7}$$

These infinitesimal symmetries will generate a Lie algebra $\mathfrak{g}$ over the field of real or complex numbers. The commutator table of Lie algebra $\mathfrak{g}$ for (1) is given below, where the entry in the i th row and j th column is defined as $[X_i, X_j] = X_i.X_j - X_j.X_i$, $i, j = 1, \dots, 13$.

	X_1	X_2	X_3	X_4	X_5	X_6	X_7	X_8	X_9	X_{10}	X_{11}	X_{12}	X_{13}
X_1	0	0	0	0	0	0	0	0	X_1	$-X_2$	0	X_3	0
X_2	0	0	0	0	0	0	0	0	X_2	X_1	X_3	0	0
X_3	0	0	0	0	0	0	0	0	X_3	0	$-X_2$	$-X_1$	0
X_4	0	0	0	0	X_1	X_2	X_3	X_4	0	0	0	0	0
X_5	0	0	0	$-X_1$	0	0	0	$-X_5$	X_5	$-X_6$	0	X_7	0
X_6	0	0	0	$-X_2$	0	0	0	$-X_6$	X_6	X_5	X_7	0	0
X_7	0	0	0	$-X_3$	0	0	0	$-X_7$	X_7	0	$-X_6$	$-X_5$	0
X_8	0	0	0	$-X_4$	X_5	X_6	X_7	0	0	0	0	0	0
X_9	$-X_1$	$-X_2$	$-X_3$	0	$-X_5$	$-X_6$	$-X_7$	0	0	0	0	0	0
X_{10}	X_2	$-X_1$	0	0	X_6	$-X_5$	0	0	0	0	$-X_{12}$	X_{11}	0
X_{11}	0	$-X_3$	X_2	0	0	$-X_7$	X_6	0	0	X_{12}	0	$-X_{10}$	0
X_{12}	$-X_3$	0	X_1	0	$-X_7$	0	X_5	0	0	$-X_{11}$	X_{10}	0	0
X_{13}	0	0	0	0	0	0	0	0	0	0	0	0	0

3 Reduction of the system

Now, we look at the group-invariant solutions to the system (1).

The system (1) is a sub-manifold M of jet space $J^2(\mathbb{R}^4, \mathbb{R}^5)$ defined by the following equations

$$u_t + uu_x + vu_y + wu_z + p_x/q = 0,$$

$$\begin{aligned}
v_t + uv_x + vv_y + wv_z + p_y/q &= 0, \\
w_t + uw_x + vw_y + ww_z + p_z/q &= 0, \\
q_t + q(u_x + v_y + w_z) + uq_x + vq_y + wq_z &= 0, \\
p_t + \gamma p(u_x + v_y + w_z) + up_x + vp_y + wp_z &= 0.
\end{aligned} \tag{8}$$

Doing as section 3.1 of [6] and find (in a sense) the most general group-invariant solutions to the system (1).

Theorem 2. *The one-parameter groups $g_i(s) : M \rightarrow M$ generated by the X_i , $i = 1, \dots, 13$ are given in the following table:*

$$\begin{aligned}
g_1 &: (t, x + s, y, z, u, v, w, p, q) \\
g_2 &: (t, x, y + s, z, u, v, w, p, q) \\
g_3 &: (t, x, y, z + s, u, v, w, p, q) \\
g_4 &: (t + s, x, y, z, u, v, w, p, q) \\
g_5 &: (t, x + st, y, z, u + s, v, w, p, q) \\
g_6 &: (t, x, y + st, z, u, v + s, w, p, q) \\
g_7 &: (t, x, y, z + st, u, v, w + s, p, q) \\
g_8 &: (e^s t, x, y, z, e^{-s} u, e^{-s} v, e^{-s} w, p, e^{2s} q) \\
g_9 &: (t, e^s x, e^s y, e^s z, e^s u, e^s v, e^s w, p, e^{-2s} q) \\
g_{10} &: (t, x \cos s + y \sin s, y \cos s - x \sin s, z, \\
&\quad v \sin s + u \cos s, v \cos s - u \sin s, w, p, q) \\
g_{11} &: (t, x, y \cos s - z \sin s, z \cos s + y \sin s, \\
&\quad u, v \cos s - w \sin s, w \cos s + v \sin s, p, q) \\
g_{12} &: (t, x \cos s - z \sin s, y, x \sin s + z \cos s, \\
&\quad u \cos s - w \sin s, v, w \cos s + u \sin s, p, q) \\
g_{13} &: (t, x, y, z, u, v, w, e^s p, e^s q)
\end{aligned} \tag{9}$$

Where, entries give the transformed point

$$\exp(sX_i)(t, x, y, z, u, v, w, p, q) = (\tilde{t}, \tilde{x}, \tilde{y}, \tilde{z}, \tilde{u}, \tilde{v}, \tilde{w}, \tilde{p}, \tilde{q}).$$

Since each group g_i , $i = 1, \dots, 13$, is a symmetry group of (1), then

Theorem 3. *Let $i = 1, \dots, 13$, $s \in \mathbb{R}$, and $\lambda \in \mathbb{R} - \{0\}$. If*

$$\begin{aligned}
u &= U(t, x, y, z), \quad v = V(t, x, y, z), \quad w = W(t, x, y, z), \\
p &= P(t, x, y, z), \quad q = Q(t, x, y, z),
\end{aligned}$$

is a solution of the system (1), so are the functions

$$\begin{aligned} u^{(i)} &= U^{(i)}(t, x, y, z), \quad v^{(i)} = V^{(i)}(t, x, y, z), \quad w^{(i)} = W^{(i)}(t, x, y, z), \\ p^{(i)} &= P^{(i)}(t, x, y, z), \quad q^{(i)} = Q^{(i)}(t, x, y, z), \end{aligned}$$

where

$$\begin{aligned} u^{(1)} &= U(t, x + s, y, z), & u^{(2)} &= U(t, x, y + s, z), \\ v^{(1)} &= V(t, x + s, y, z), & v^{(2)} &= V(t, x, y + s, z), \\ w^{(1)} &= W(t, x + s, y, z), & w^{(2)} &= W(t, x, y + s, z), \\ p^{(1)} &= P(t, x + s, y, z), & p^{(2)} &= P(t, x, y + s, z), \\ q^{(1)} &= Q(t, x + s, y, z), & q^{(2)} &= Q(t, x, y + s, z). \end{aligned}$$

$$\begin{aligned} u^{(3)} &= U(t, x, y, z + s), & u^{(4)} &= U(t + s, x, y, z), \\ v^{(3)} &= V(t, x, y, z + s), & v^{(4)} &= V(t + s, x, y, z), \\ w^{(3)} &= W(t, x, y, z + s), & w^{(4)} &= W(t + s, x, y, z), \\ p^{(3)} &= P(t, x, y, z + s), & p^{(4)} &= P(t + s, x, y, z), \\ q^{(3)} &= Q(t, x, y, z + s), & q^{(4)} &= Q(t + s, x, y, z). \end{aligned}$$

$$\begin{aligned} u^{(5)} &= U(t, x + st, y, z) - s, & u^{(6)} &= U(t, x, y + st, z), \\ v^{(5)} &= V(t, x + st, y, z), & v^{(6)} &= V(t, x, y + st, z) - s, \\ w^{(5)} &= W(t, x + st, y, z), & w^{(6)} &= W(t, x, y + st, z), \\ p^{(5)} &= P(t, x + st, y, z), & p^{(6)} &= P(t, x, y + st, z), \\ q^{(5)} &= Q(t, x + st, y, z), & q^{(6)} &= Q(t, x, y + st, z). \end{aligned}$$

$$\begin{aligned} u^{(7)} &= U(t, x, y, z + st), & u^{(8)} &= \lambda.U(\lambda.t, x, y, z), \\ v^{(7)} &= V(t, x, y, z + st), & v^{(8)} &= \lambda.V(\lambda.t, x, y, z), \\ w^{(7)} &= W(t, x, y, z + st) - s, & w^{(8)} &= \lambda.W(\lambda.t, x, y, z), \\ p^{(7)} &= P(t, x, y, z + st), & p^{(8)} &= P(\lambda.t, x, y, z), \\ q^{(7)} &= Q(t, x, y, z + st), & q^{(8)} &= \frac{1}{\lambda^2}.Q(\lambda.t, x, y, z). \end{aligned}$$

$$\begin{aligned} u^{(9)} &= \frac{1}{\lambda}.U(t, \lambda.x, \lambda.y, \lambda.z), & u^{(13)} &= U(t, x, y, z), \\ v^{(9)} &= \frac{1}{\lambda}.V(t, \lambda.x, \lambda.y, \lambda.z), & v^{(13)} &= V(t, x, y, z), \\ w^{(9)} &= \frac{1}{\lambda}.W(t, \lambda.x, \lambda.y, \lambda.z), & w^{(13)} &= W(t, x, y, z), \\ p^{(9)} &= P(t, \lambda.x, \lambda.y, \lambda.z), & p^{(13)} &= \lambda.P(t, x, y, z), \\ q^{(9)} &= \lambda^2.Q(t, \lambda.x, \lambda.y, \lambda.z), & q^{(13)} &= \lambda.Q(t, x, y, z). \end{aligned}$$

$$\begin{aligned} u^{(10)} &= \cos s.U(t, x \cos s + y \sin s, y \cos s - x \sin s, z) \\ &\quad - \sin s.V(t, x \cos s + y \sin s, y \cos s - x \sin s, z), \\ v^{(10)} &= \sin s.U(t, x \cos s + y \sin s, y \cos s - x \sin s, z) \\ &\quad + \cos s.V(t, x \cos s + y \sin s, y \cos s - x \sin s, z), \end{aligned}$$

$$\begin{aligned}
w^{(10)} &= W(t, x \cos s + y \sin s, y \cos s - x \sin s, z), \\
p^{(10)} &= P(t, x \cos s + y \sin s, y \cos s - x \sin s, z), \\
q^{(10)} &= Q(t, x \cos s + y \sin s, y \cos s - x \sin s, z). \\
\\
u^{(11)} &= U(t, x, y \cos s - z \sin s, z \cos s + y \sin s), \\
v^{(11)} &= \cos s.V(t, x, y \cos s - z \sin s, z \cos s + y \sin s) \\
&\quad + \sin s.W(t, x, y \cos s - z \sin s, z \cos s + y \sin s), \\
w^{(11)} &= \cos s.W(t, x, y \cos s - z \sin s, z \cos s + y \sin s) \\
&\quad - \sin s.V(t, x, y \cos s - z \sin s, z \cos s + y \sin s), \\
p^{(11)} &= P(t, x, y \cos s - z \sin s, z \cos s + y \sin s), \\
q^{(11)} &= Q(t, x, y \cos s - z \sin s, z \cos s + y \sin s). \\
\\
u^{(12)} &= \cos s.U(t, x \cos s - z \sin s, y, x \sin s + z \cos s) \\
&\quad + \sin s.W(t, x \cos s - z \sin s, y, x \sin s + z \cos s), \\
v^{(12)} &= V(t, x \cos s - z \sin s, y, x \sin s + z \cos s), \\
w^{(12)} &= \cos s.W(t, x \cos s - z \sin s, y, x \sin s + z \cos s) \\
&\quad - \sin s.U(t, x \cos s - z \sin s, y, x \sin s + z \cos s), \\
p^{(12)} &= P(t, x \cos s - z \sin s, y, x \sin s + z \cos s), \\
q^{(12)} &= Q(t, x \cos s - z \sin s, y, x \sin s + z \cos s).
\end{aligned}$$

4 Structure of Lie algebra $\mathfrak{g}$

In this section, we determine the structure of full symmetry algebra $\mathfrak{g}$. of system (1).

The center $\mathfrak{z}$ of $\mathfrak{g}$ is $\text{Span}_{\mathbb{R}}\{X_{13}\}$. Therefore, the quotient algebra $\mathfrak{g}_1 := \mathfrak{g}/\mathfrak{z}$ is $\text{Span}_{\mathbb{R}}\{Y_1, \dots, Y_{12}\}$; where $Y_i := X_i + \mathfrak{z}$, $i = 1, \dots, 12$. The commutator table of Lie algebra $\mathfrak{g}_1$ is given below, where the entry in the i th row and j th column is defined as $[Y_i, Y_j] = Y_i.Y_j - Y_j.Y_i$,

$i, j = 1, \dots, 12$.

	Y_1	Y_2	Y_3	Y_4	Y_5	Y_6	Y_7	Y_8	Y_9	Y_{10}	Y_{11}	Y_{12}
Y_1	0	0	0	0	0	0	0	0	Y_1	$-Y_2$	0	Y_3
Y_2	0	0	0	0	0	0	0	0	Y_2	Y_1	Y_3	0
Y_3	0	0	0	0	0	0	0	0	Y_3	0	$-Y_2$	$-Y_1$
Y_4	0	0	0	0	Y_1	Y_2	Y_3	Y_4	0	0	0	0
Y_5	0	0	0	$-Y_1$	0	0	0	$-Y_5$	Y_5	$-Y_6$	0	Y_7
Y_6	0	0	0	$-Y_2$	0	0	0	$-Y_6$	Y_6	Y_5	Y_7	0
Y_7	0	0	0	$-Y_3$	0	0	0	$-Y_7$	Y_7	0	$-Y_6$	$-Y_5$
Y_8	0	0	0	$-Y_4$	Y_5	Y_6	Y_7	0	0	0	0	0
Y_9	$-Y_1$	$-Y_2$	$-Y_3$	0	$-Y_5$	$-Y_6$	$-Y_7$	0	0	0	0	0
Y_{10}	Y_2	$-Y_1$	0	0	Y_6	$-Y_5$	0	0	0	0	$-Y_{12}$	Y_{11}
Y_{11}	0	$-Y_3$	Y_2	0	0	$-Y_7$	Y_6	0	0	Y_{12}	0	$-Y_{10}$
Y_{12}	$-Y_3$	0	Y_1	0	$-Y_7$	0	Y_5	0	0	$-Y_{11}$	Y_{10}	0

The algebra $\mathfrak{g}_1$ is not solvable, because

$$\mathfrak{g}_1^{(1)} := [\mathfrak{g}_1, \mathfrak{g}_1] = \text{Span}_{\mathbb{R}}\{Y_1, \dots, Y_{10}\}, \text{ and } \mathfrak{g}_1^{(2)} := [\mathfrak{g}_1^{(1)}, \mathfrak{g}_1^{(1)}] = \mathfrak{g}_1^{(1)}.$$

Lie algebra $\mathfrak{g}_1$ admits a Levi-decomposition into the semi-direct sum

$$\mathfrak{g}_1 = \mathfrak{r} \ltimes \mathfrak{s},$$

where $\mathfrak{r} = \text{Span}_{\mathbb{R}}\{Y_1, \dots, Y_9\}$ is the radical of $\mathfrak{g}_1$ (the largest solvable ideal contained in $\mathfrak{g}_1$) and $\mathfrak{s} = \text{Span}_{\mathbb{R}}\{Y_{10}, Y_{11}, Y_{12}\}$ is a semi-simple subalgebra of $\mathfrak{g}_1$. The radical $\mathfrak{r}$ is uniquely defined but the semi-simple subalgebra $\mathfrak{s}$ is not. $\mathfrak{s}$ as an algebra is isomorphic to 3-dimensional special orthogonal algebra $\mathfrak{so}(3) := \{A \in \text{Mat}(3 \times 3; \mathbb{R}) \mid A^t = -A\}$, which is a simple algebra.

The radical $\mathfrak{r}$ is solvable, with and

$$\mathfrak{r}^{(1)} = \text{Span}_{\mathbb{R}}\{Y_1, \dots, Y_7\}, \quad \mathfrak{r}^{(2)} = \text{Span}_{\mathbb{R}}\{Y_1, Y_2, Y_3\}, \quad \mathfrak{r}^{(3)} = \{0\}.$$

It is semi-direct sum $\mathfrak{r} = \mathfrak{r}^{(1)} \ltimes \mathfrak{g}_2$, where $\mathfrak{g}_2 = \text{Span}_{\mathbb{R}}\{Y_8, Y_9\}$ is isomorphic to the two dimensional Abelian Lie algebra $\mathbb{R}^2$; i.e.

$$\mathfrak{r} = \mathfrak{r}^{(1)} \ltimes \mathbb{R}^2.$$

$\mathfrak{r}^{(1)}$ is semi-direct sum $\mathfrak{r}^{(1)} = \mathfrak{r}^{(2)} \ltimes \mathfrak{g}_3$, where $\mathfrak{g}_3 = \text{Span}_{\mathbb{R}}\{Y_4, \dots, Y_7\}$ is isomorphic to the four dimensional Abelian Lie algebra $\mathbb{R}^4$, and $\mathfrak{r}^{(2)}$ is isomorphic to the three dimensional Abelian Lie algebra $\mathbb{R}^3$; i.e.

$$\mathfrak{r}^{(1)} = \mathbb{R}^3 \ltimes \mathbb{R}^4.$$

Theorem 4. *The symmetry algebra $\mathfrak{g}$ of system (1) have the following structure*

$$\mathfrak{g} \cong \mathbb{R} \times ((\mathbb{R}^3 \ltimes \mathbb{R}^4) \ltimes \mathbb{R}^2) \ltimes \mathfrak{so}(3).$$

Theorem 5. *There is a normal Lie-subalgebra of symmetry Lie-group of system (1) which is isomorphic to the Galilean group of $\mathbb{R}^4$:*

$$\text{Gal}(4, \mathbb{R}) = \left\{ \begin{pmatrix} 1 & \mathbf{0} & s \\ \mathbf{v} & \mathbf{R} & \mathbf{y} \\ 0 & \mathbf{0} & 1 \end{pmatrix} \mid \mathbf{R} \in \text{O}(3, \mathbb{R}), s \in \mathbb{R}, \text{ and } \mathbf{y}, \mathbf{v} \in \mathbb{R}^3 \right\}.$$

Therefore, the system (1) is invariant up to Galilean motions of space-times $\mathbb{R}^4$ (see [3]).

Proof. Let $\mathfrak{b} = \text{Span}_{\mathbb{R}}\{X_1, \dots, X_7, X_{10}, X_{11}, X_{12}\}$. Lie algebra structure of $\mathfrak{b}$ is as $\mathfrak{gal}(3)$, the lie algebra of Galilean group $\text{Gal}(4, \mathbb{R})$, and $\mathfrak{b}$ an ideal of $\mathfrak{g}$. Therefore, there is a Lie-subgroup of G such that its Lie-algebra is $\mathfrak{b}$, by the Theorem 2.53 of Olver [6]. $\square$

Conclusion 1. $\mathfrak{g}$ is semi-direct sum of

$$\mathfrak{gal}(4) \cong \text{Span}_{\mathbb{R}}\{X_1, \dots, X_7, X_{10}, X_{11}, X_{12}\}$$

and 3-dimensional Abelian algebra $\mathbb{R}^3 \cong \text{Span}\{X_8, X_9, X_{13}\}$.

5 Optimal system of sub-algebras

As is well known, the Lie group theoretic method plays an important role in finding exact solutions and performing symmetry reductions of differential equations. Since any linear combination of infinitesimal generators is also an infinitesimal generator, there are always infinitely many different symmetry subgroups for the differential equation. So, a mean of determining which subgroups would give essentially different types of solutions is necessary and significant for a complete understanding of the invariant solutions. As any transformation in the full symmetry group maps a solution to another solution, it is sufficient to find invariant solutions which are not related by transformations in the full symmetry group, this has led to the concept of an optimal system [7]. The problem of finding an optimal system of subgroups is equivalent to that of finding an optimal system of subalgebras. For one-dimensional subalgebras, this classification problem is essentially the same as the problem of classifying the orbits of the adjoint representation. This problem is attacked by the naive approach of taking a general element in the Lie algebra and subjecting it to various adjoint transformations so as to simplify it as much as possible. The idea of using the adjoint representation to classify group-invariant solutions was due to [7] and [5].

The adjoint action is given by the Lie series

$$\text{Ad}(\exp(sX_i)X_j) = X_j - s[X_i, X_j] + \frac{s^2}{2}[X_i, [X_i, X_j]] - \dots, \quad (10)$$

where $[X_i, X_j]$ is the commutator for the Lie algebra, s is a parameter, and $i, j = 1, \dots, 13$. We can write the adjoint action for the Lie algebra $\mathfrak{g}$, and show that

Theorem 6. *A one-dimensional optimal system of (1) is given by*

- 1) $X = X_8 + \sum_{i=9}^{13} a_i X_i,$
- 2) $X = a_7 X_7 + X_8 + X_9 + \sum_{i=10}^{13} a_i X_i,$
- 3) $X = a_6 X_6 + a_7 X_7 + X_8 + X_9 + \sum_{i=11}^{13} a_i X_i,$
- 4) $X = a_5 X_5 + X_8 + X_9 + a_{11} X_i + a_{13} X_{13},$
- 5) $X = \sum_{i=5}^7 a_i X_i + X_8 + X_9 + a_{13} X_{13},$
- 6) $X = a_4 X_4 + X_9 + \sum_{i=10}^{13} a_i X_i,$
- 7) $X = X_4 + a_5 X_5 + a_8 X_8 + \sum_{i=10}^{13} a_i X_i,$
- 8) $X = X_4 + a_6 X_6 + a_8 X_8 + a_{10} X_{10} + a_{12} X_{12} + a_{13} X_{13},$
- 9) $X = X_4 + a_7 X_7 + a_8 X_8 + a_{10} X_{10} + a_{12} X_{12} + a_{13} X_{13},$
- 10) $X = X_4 + \sum_{i=5}^8 a_i X_i + a_{13} X_{13},$
- 11) $X = X_4 + a_7 X_7 + \sum_{i=10}^{13} a_i X_i,$
- 12) $X = X_7 + \sum_{i=10}^{13} a_i X_i,$
- 13) $X = a_3 X_3 + \sum_{i=10}^{13} a_i X_i,$
- 14) $X = X_4 + a_6 X_6 + \sum_{i=11}^{13} a_i X_i,$
- 15) $X = X_4 + a_5 X_5 + a_{11} X_{11} + a_{13} X_{13},$
- 16) $X = X_4 + \sum_{i=5}^7 a_i X_i + a_{13} X_{13},$
- 17) $X = X_6 + \sum_{i=11}^{13} a_i X_i,$
- 18) $X = a_2 X_2 + \sum_{i=11}^{13} a_i X_i,$
- 19) $X = a_2 X_2 + a_{12} X_{12} + a_{13} X_{13},$
- 20) $X = X_5 + a_{11} X_{11} + a_{13} X_{13},$
- 21) $X = a_2 X_2 + a_3 X_3 + X_5 + a_{13} X_{13},$
- 22) $X = a_1 X_1 + X_{11} + a_{13} X_{13},$
- 23) $X = a_1 X_1 + a_3 X_3 + X_6 + a_{13} X_{13},$
- 24) $X = a_1 X_1 + a_2 X_2 + a_3 X_3 + a_{13} X_{13},$
- 25) $X = X_5 + a_{10} X_{10} + a_{11} X_{11} + a_{13} X_{13},$
- 26) $X = a_3 X_3 + X_{10} + a_{11} X_{11} + a_{13} X_{13}.$

Proof: $F(sX_i) : \mathfrak{g} \rightarrow \mathfrak{g}$ defined by $X \mapsto \text{Ad}(\exp(sX_i)X)$ is a linear map, for $i = 1, \dots, 13$. The matrix M_{sX_i} of $F(sX_i)$, $i = 1, \dots, 13$, with respect to basis $\{X_1, \dots, X_{13}\}$ is

$$\begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ -s & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & s & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & -s & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix},$$

[illegible]

[illegible]

11

[illegible]

[illegible]

[illegible]

[illegible]

[illegible]

$$\begin{bmatrix} \cos s & 0 & \sin s & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ -\sin s & 0 & \cos s & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & \cos s & 0 & \sin s & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & -\sin s & 0 & \cos s & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & \cos s & \sin s & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & -\sin s & \cos s & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix},$$

and I_{13} , respectively. Let $X = \sum_{i=1}^{13} a_i X_i$, then

$$\begin{aligned} F(s_7.X_7) \circ F(s_6.X_6) \circ \cdots \circ F(s_1.X_1) : X \mapsto \\ (s_3a_{12} - s_2a_{10} - s_1a_9 + a_1 - s_5s_4a_8 + s_5a_4 - s_4a_5).X_1 \\ + (-s_6s_4a_8 + s_3a_{11} - s_2a_9 + s_1a_{10} + a_2 + s_6a_4 - s_4a_6).X_2 \\ + (-s_7s_4a_8 - s_3a_9 - s_2a_{11} - s_1a_{12} + a_3 + s_7a_4 - s_4a_7).X_3 \quad (11) \\ + (a_4 - s_4a_8).X_4 + (-s_6a_{10} - s_5a_9 + s_5a_8 + a_5 + s_7a_{12}).X_5 \\ + (-s_6a_9 + s_6a_8 + s_5a_{10} + a_6 + s_7a_{11}).X_6 \\ + (s_7a_8 - s_6a_{11} - s_5a_{12} + a_7 - s_7a_9).X_7 \\ + a_8.X_8 + a_9.X_9 + a_{10}.X_{10} + a_{11}.X_{11} + a_{12}.X_{12} + a_{13}.X_{13} \end{aligned}$$

If a_8 , and $a_9 \neq 0$, and $a_8 \neq 2a_9$, then we can make the coefficients of $X_1, \dots, X_7$ vanish. Scaling X if necessary, we can assume that $a_8 = 1$. And X is reduced to Case 1.

If $a_8 = a_9 \neq 0$, and $a_{10} \neq 0$, then we can make the coefficients of $X_1, \dots, X_6$ vanish. Scaling X if necessary, we can assume that $a_8 = 1$. And X is reduced to Case 2.

If $a_8 = a_9 \neq 0$, $a_{10} = 0$, and $a_{12} \neq 0$, then we can make the coefficients of $X_1, \dots, X_5$, and X_{10} vanish. Scaling X if necessary, we can assume that $a_8 = 1$. And X is reduced to Case 3.

If $a_8 = a_9 \neq 0$, $a_{10} = a_{12} = 0$, and $a_{11} \neq 0$, then we can make the coefficients of $X_1, \dots, X_4$, X_6 , X_7 , X_{10} , and X_{12} vanish. Scaling X if necessary, we can assume that $a_8 = 1$. And X is reduced to Case 4.

If $a_8 = a_9 \neq 0$, $a_{10} = a_{11} = a_{12} = 0$, then we can make the coefficients of $X_1, \dots, X_4$, and $X_{10}, \dots, X_{11}$ vanish. Scaling X if necessary, we can assume that $a_8 = 1$. And X is reduced to Case 5.

If $a_8 = 0$, and $a_9 \neq 0$, then we can make the coefficients of $X_1, \dots, X_3$, and $X_5, \dots, X_8$ vanish. Scaling X if necessary, we can assume that $a_9 = 1$. And X is reduced to Case 6.

If $a_8 \neq 0$, $a_9 = 0$, and $a_{11} \neq 0$, then we can make the coefficients of $X_1, \dots, X_3$, and X_6 , X_7 , and X_9 vanish. And X is reduced to Case 7.

If $a_8 \neq 0$, $a_9 = a_{11} = 0$, and $a_{12} \neq 0$, then we can make the coefficients of $X_1, \dots, X_3$, and X_5 , X_7 , X_9 , and X_{11} vanish. And X is reduced to Case 8.

If $a_8 \neq 0$, $a_9 = a_{11} = a_{12} = 0$, and $a_{10} \neq 0$, then we can make the coefficients of $X_1, \dots, X_3$, and X_5 , X_6 , X_9 , and X_{11} vanish. And X is reduced to Case 9.

If $a_8 \neq 0$, and $a_9 = a_{10} = a_{11} = a_{12} = 0$, then we can make the coefficients of $X_1, \dots, X_3$, and $X_9, \dots, X_{12}$ vanish. And X is reduced to Case 10.

If $a_8 = a_9 = 0$, $a_4 \neq 0$, and $a_{10} \neq 0$, then we can make the coefficients of $X_1, \dots, X_3$, X_5 , X_6 , X_8 , and X_9 vanish. Scaling X if necessary, we can assume that $a_4 = 1$. And X is reduced to Case 11.

If $a_4 = a_8 = a_9 = 0$, $a_{10} \neq 0$, $a_{12} \neq 0$, and $a_6 a_{12} + a_7 a_{10} \neq a_5 a_{11}$, then we can make the coefficients of $X_1, \dots, X_6$, X_8 , and X_9 vanish. Scaling X if necessary, we can assume that $a_7 = 1$. And X is reduced to Case 12.

If $a_4 = a_8 = a_9 = 0$, $a_{10} \neq 0$, $a_{12} \neq 0$, and $a_6 a_{12} + a_7 a_{10} = a_5 a_{11}$, then we can make the coefficients of X_1 , X_2 , and $X_4, \dots, X_9$ vanish. And X is reduced to Case 13.

If $a_8 = a_9 = a_{10} = 0$, and $a_4 \neq 0$, then we can make the coefficients of $X_1, \dots, X_3$, X_5 , and $X_7, \dots, X_{10}$ vanish. Scaling X if necessary, we can assume that $a_4 = 1$. And X is reduced to Case 14.

If $a_8 = \dots = a_{10} = 0$, $a_{12} = 0$, $a_4 \neq 0$, and $a_{11} \neq 0$, then we can make the coefficients of $X_1, \dots, X_3$, and $X_8, \dots, X_{10}$ vanish. Scaling X if necessary, we can assume that $a_4 = 1$. And X is reduced to Case 15.

If $a_8 = \dots = a_{12} = 0$, and $a_4 \neq 0$, then we can make the coefficients of $X_1, \dots, X_3$, and $X_8, \dots, X_{10}$ vanish. Scaling X if necessary, we can assume that $a_4 = 1$. And X is reduced to Case 16.

If $a_4 = a_8 = a_9 = a_{10} = 0$, $a_{12} \neq 0$, and $a_5 a_{11} \neq a_6 a_{12}$, then we can make the coefficients of $X_1, \dots, X_5$, and $X_7, \dots, X_{10}$ vanish. Scaling X if necessary, we can assume that $a_6 = 1$. And X is reduced to Case 17.

If $a_4 = a_8 = a_9 = a_{10} = 0$, $a_{12} \neq 0$, $a_5 a_{11} = a_6 a_{12}$, and $a_6 \neq 0$, then we can make the coefficients of $X_1, \dots, X_5$, and $X_7, \dots, X_{10}$ vanish. And X is reduced to Case 18.

If $a_4 = a_6 = a_8 = a_9 = a_{10} = 0$, $a_{12} \neq 0$, $a_5 a_{11} = a_6 a_{12}$, and $a_6 \neq 0$, then we can make the coefficients of $X_1, \dots, X_5$, and $X_7, \dots, X_{10}$ vanish. And X is reduced to Case 18.

If $a_4 = a_6 = a_8 = \dots = a_{11} = 0$, $a_{12} \neq 0$, $a_5 a_{11} = a_6 a_{12}$, $a_6 \neq 0$, and $a_5 \neq 0$, then we can make the coefficients of X_1 , and $X_3, \dots, X_{11}$ vanish. And X is reduced to Case 19.

If $a_4 = a_8 = a_9 = a_{10} = a_{12} = 0$, $a_5 \neq 0$, and $a_{11} \neq 0$, then we can make the coefficients of $X_1, \dots, a_4$, $X_6, \dots, X_{10}$, and a_{12} vanish. Scaling X if necessary, we can assume that $a_5 = 1$. And X is reduced to Case 20.

If $a_4 = a_8 = \dots = a_{12} = 0$, and $a_5 \neq 0$, then we can make the coefficients of X_1 , a_4 , and $X_8, \dots, X_{12}$ vanish. Scaling X if necessary, we can assume that $a_5 = 1$. And X is reduced to Case 21.

If $a_4 = a_5 = a_8 = a_9 = a_{10} = a_{12} = 0$, and $a_{11} \neq 0$, then we can make the coefficients of $X_2, \dots, X_{10}$ and a_{12} vanish. Scaling X if necessary, we can assume that $a_{11} = 1$. And X is reduced to Case 22.

If $a_4 = a_5 = a_8 = \dots = a_{12} = 0$, and $a_6 \neq 0$, then we can make the coefficients of X_2 , a_5 , and $X_7, \dots, X_{12}$ vanish. Scaling X if necessary, we can assume that $a_6 = 1$. And X is reduced to Case 23.

If $a_4 = a_5 = a_6 = 0$, and $a_8 = \dots = a_{12} = 0$, then we can make the coefficients of $X_4, \dots, X_{12}$ vanish. And X is reduced to Case 24.

If $a_4 = a_8 = a_9 = a_{12} = 0$, $a_{10} \neq 0$, and $a_7 a_{10} \neq a_5 a_{11}$, then we can make the coefficients of $X_1, \dots, X_4$, $X_6, \dots, X_9$, and $a_{12} = 0$ vanish. Scaling X if necessary, we can assume that $a_5 = 1$. And X is reduced to Case 25.

If $a_4 = a_8 = a_9 = a_{12} = 0$, $a_7 a_{10} = a_5 a_{11}$, and $a_{10} \neq 0$, then we can make the coefficients of $X_1, \dots, X_4$, $X_6, \dots, X_9$, and $a_{12} = 0$ vanish. Scaling X if necessary, we can assume that $a_{10} = 1$. And X is reduced to Case 26. And the Theorem follows. $\square$

According to our optimal system of one-dimensional subalgebras of the full symmetry algebra $\mathfrak{g}$, we need only find group-invariant solutions for 26 one-parameter subgroups generated by X as Theorem 6. For example, as a direct consequence of Theorem 3 and case 1 of Theorem 6, we deduce the following conclusion:

Conclusion 2 *Let $s, c_1, \dots, c_6 \in \mathbb{R}$ with c_1, c_2 , and $c_6 \neq 0$, and s be sufficiently small. Then, if*

$$\begin{aligned} u &= U(t, x, y, z), \quad v = V(t, x, y, z), \quad w = W(t, x, y, z), \\ p &= P(t, x, y, z), \quad q = Q(t, x, y, z), \end{aligned}$$

is a solution of the system (1), so are the functions

$$\begin{aligned} \tilde{u} &= c_1^s c_2^{-s} (\cos(c_3 s) \cos(c_5 s) - \sin(c_3 s) \sin(c_4 s) \sin(c_5 s)) \cdot U(\tilde{t}, \tilde{x}, \tilde{y}, \tilde{z}) \\ &\quad - c_1^s c_2^{-s} (\sin(c_3 s) \cos(c_5 s) + \cos(c_3 s) \sin(c_4 s) \cos(c_5 s)) \cdot V(\tilde{t}, \tilde{x}, \tilde{y}, \tilde{z}) \\ &\quad + c_1^s c_2^{-s} \cos(c_4 s) \sin(c_5 s) \cdot W(\tilde{t}, \tilde{x}, \tilde{y}, \tilde{z}), \\ \tilde{v} &= c_1^s c_2^{-s} \sin(c_3 s) \cos(c_4 s) \cdot U(\tilde{t}, \tilde{x}, \tilde{y}, \tilde{z}) \\ &\quad + c_1^s c_2^{-s} \cos(c_3 s) \cos(c_4 s) \cdot V(\tilde{t}, \tilde{x}, \tilde{y}, \tilde{z}) \\ &\quad + c_1^s c_2^{-s} \cdot \sin(c_4 s) \cdot W(\tilde{t}, \tilde{x}, \tilde{y}, \tilde{z}), \\ \tilde{w} &= -c_1^s c_2^{-s} (\cos(c_3 s) \sin(c_5 s) + \sin(c_3 s) \sin(c_4 s) \cos(c_5 s)) \cdot U(\tilde{t}, \tilde{x}, \tilde{y}, \tilde{z}) \\ &\quad + c_1^s c_2^{-s} (\sin(c_3 s) \sin(c_5 s) - \cos(c_3 s) \sin(c_4 s) \cos(c_5 s)) \cdot V(\tilde{t}, \tilde{x}, \tilde{y}, \tilde{z}) \\ &\quad + c_1^s c_2^{-s} \cos(c_4 s) \cos(c_5 s) \cdot W(\tilde{t}, \tilde{x}, \tilde{y}, \tilde{z}), \\ \tilde{p} &= c_6^{-s} \cdot P(\tilde{t}, \tilde{x}, \tilde{y}, \tilde{z}), \\ \tilde{q} &= c_1^{-2s} c_2^{2s} c_6^{-s} \cdot Q(\tilde{t}, \tilde{x}, \tilde{y}, \tilde{z}), \end{aligned}$$

where $\tilde{t} = c_1^s t$, and

$$\begin{aligned} \tilde{x} &= c_2^s (\sin(c_5 s) \sin(c_4 s) \sin(c_3 s) + \cos(c_5 s) \cos(c_3 s)) \cdot x \\ &\quad + c_2^s (\sin(c_3 s) \cos(c_5 s) - \cos(c_3 s) \sin(c_4 s) \sin(c_5 s)) \cdot y \\ &\quad - c_2^s \cos(c_4 s) \sin(c_5 s) \cdot z, \\ \tilde{y} &= -c_2^s \sin(c_3 s) \cos(c_4 s) \cdot x + (c_2 s) \cos(c_3 s) \cos(c_4 s) \cdot y - c_2^s \sin(c_4 s) \cdot z, \\ \tilde{z} &= c_2^s (\cos(c_3 s) \sin(c_5 s) - \sin(c_3 s) \sin(c_4 s) \cos(c_5 s)) \cdot x \end{aligned}$$

$$\begin{aligned}
&+c_2^s(\cos(c_3s)\sin(c_4s)\cos(c_5s)+\sin(c_3s)\sin(c_5s)).y \\
&+c_2^s\cos(c_4s)\cos(c_5s).z.
\end{aligned}$$

The construction of the group-invariant solutions for each of the one-dimensional subgroups in the optimal system proceeds in the same fashion.

6 Acknowledgements

I am grateful to professor Ian M. Anderson for his consideration and his mathematical career to the development the "DifferentialGeometry" package of MAPLE 11. It is an applicable tool to extremely complex and tedious geometrical computations.

References

- [1] G.W. BLUMAN, J.D. COLE, *Similarity Methods for Differential Equations*, Springer, Berlin, 1974.
- [2] N.H. IBRAGIMOV, *Transformation Groups Applied to Mathematical Physics*, Reidel, Dordrecht, 1985.
- [3] M. NADJAFIKHAH, A. MAHDIPOUR-SHIRAYEH, *Galilean Classification of Spacetime Curves*, arXiv e-print: arXiv:0711.1979.
- [4] F. OLIVERI, M.P. SPECIALE, *Exact solutions to the unsteady equations of perfect gases through Lie group analysis and substitution principles*, Int. J. Non-Linear Mech. 37 (2002) 257–274.
- [5] P.J. OLVER, *Applications of Lie Groups to Differential Equations*, Springer, New York, 1986.
- [6] P.J. OLVER, *Equivalence, Invariants, and Symmetry*, Cambridge University Press, 1995.
- [7] L.V. OVSIANNIKOV, *Group Analysis of Differential Equations*, Academic Press, New York, 1982.
- [8] T.R. SEKHAR, V.D. SHARMA, *Similarity solutions for three dimensional Euler equations using Lie group analysis*, Appl. Math. and Comp. 196 (2008) 147–157.
- [9] W.H. STEEB, *Continuous symmetries, Lie algebras, differential equations and computer algebra*, World Sci. Pub. Co., Singapore, 1996.

Group analysis of three dimensional Euler equations of gas dynamics

Mehdi Nadjafikhah^a

^a*School of Mathematics, Iran University of Science and Technology,
Narmak, Tehran, I.R.IRAN.*

Abstract

In this paper, the equations governing the unsteady flow of a perfect polytropic gas in three space dimensions are considered. The basic similarity reductions for this system are performed. Reduced equations and exact solutions associated with the symmetries are obtained. This results is used to give the invariance of system up to Galilean motions of space-times $\mathbb{R}^4$. Then, an optimal system of one-dimensional sub-algebras for symmetry algebra of this system is given.

Key words: Euler equations, Lie group of transformations, Symmetry generators, Similarity solution, optimal system of Lie sub-algebras.
2000 MSC: 35C05, 76M60.

1. Introduction

The equations governing the unsteady flow of a perfect polytropic gas in three space dimensions are

$$\begin{aligned}
 \frac{\partial u}{\partial t} + u \frac{\partial u}{\partial x} + v \frac{\partial u}{\partial y} + w \frac{\partial u}{\partial z} + \frac{1}{q} \frac{\partial p}{\partial x} &= 0, \\
 \frac{\partial v}{\partial t} + u \frac{\partial v}{\partial x} + v \frac{\partial v}{\partial y} + w \frac{\partial v}{\partial z} + \frac{1}{q} \frac{\partial p}{\partial y} &= 0, \\
 \frac{\partial w}{\partial t} + u \frac{\partial w}{\partial x} + v \frac{\partial w}{\partial y} + w \frac{\partial w}{\partial z} + \frac{1}{q} \frac{\partial p}{\partial z} &= 0, \\
 \frac{\partial q}{\partial t} + q \left(\frac{\partial u}{\partial x} + \frac{\partial v}{\partial y} + \frac{\partial w}{\partial z} \right) + u \frac{\partial q}{\partial x} + v \frac{\partial q}{\partial y} + w \frac{\partial q}{\partial z} &= 0, \\
 \frac{\partial p}{\partial t} + \gamma p \left(\frac{\partial u}{\partial x} + \frac{\partial v}{\partial y} + \frac{\partial w}{\partial z} \right) + u \frac{\partial p}{\partial x} + v \frac{\partial p}{\partial y} + w \frac{\partial p}{\partial z} &= 0.
 \end{aligned} \tag{1}$$

where t is the time and x, y, z are the space coordinates; $q(t, x, y, z)$ is the density, $p(t, x, y, z)$ the pressure, $u(t, x, y, z)$, $v(t, x, y, z)$ and $w(t, x, y, z)$ the

Email addresses: m nadjafikhah@iust.ac.ir (Mehdi Nadjafikhah)

velocity components in the x , y and z directions, respectively, and $\gamma \in \mathbb{R}$ is the adiabatic index. These equations are called "three dimensional Euler equations of gas dynamics" [8] and section 6.3 of [9].

In this paper, we consider the equations governing the unsteady flow of a perfect polytropic gas in three space dimensions. The basic similarity reductions for this system are performed. Reduced equations and exact solutions associated with the symmetries are obtained. We find an optimal system of one-dimensional sub-algebras for symmetry algebra of this system.

In section 2 we find the full symmetry algebra of system (1). Chapter 3 is devoted to the group-invariant solutions to the system. The structure of full symmetry algebra illustrated in section. The last section deals with a optimal system of sub-algebras.

2. Lie symmetry of the system

The classical method for finding symmetry reductions of PDE is the Lie group method of infinitesimal transformations. To apply the classical method to (1), we consider the one-parameter Lie group of infinitesimal transformations in (t, x, y, z, u, v, p, q) given by

$$\begin{aligned}\tilde{t} &= t + s.\xi_1(t, x, y, z, u, v, p, q) + O(s^2), \\ \tilde{x} &= x + s.\xi_2(t, x, y, z, u, v, p, q) + O(s^2), \\ \tilde{y} &= y + s.\xi_3(t, x, y, z, u, v, p, q) + O(s^2), \\ \tilde{z} &= z + s.\xi_4(t, x, y, z, u, v, p, q) + O(s^2), \\ \tilde{u} &= u + s.\eta_1(t, x, y, z, u, v, p, q) + O(s^2), \\ \tilde{v} &= v + s.\eta_2(t, x, y, z, u, v, p, q) + O(s^2), \\ \tilde{w} &= w + s.\eta_3(t, x, y, z, u, v, p, q) + O(s^2), \\ \tilde{p} &= p + s.\eta_4(t, x, y, z, u, v, p, q) + O(s^2), \\ \tilde{q} &= q + s.\eta_5(t, x, y, z, u, v, p, q) + O(s^2).\end{aligned}\tag{2}$$

where s is the group parameter. One requires that this transformation leaves the sub-manifold S invariant. Let

$$X = \xi_1 \partial_t + \xi_2 \partial_x + \xi_3 \partial_y + \xi_4 \partial_z + \eta_1 \partial_u + \eta_2 \partial_v + \eta_3 \partial_w + \eta_4 \partial_p + \eta_5 \partial_q$$

be the corresponding infinitesimal transformation; where coefficients are real-valued C^∞ -functions of $t, x, y, z, u, v, w, p, q$. The infinitesimal criterion of invariance (1) will thus involve $t, x, y, z, u, v, w, p, q$ and the derivatives of u, v, w, p, q with respect to t, x, y, z , as well as ξ_i , $i = 1, 2, 3, 4$, and η_j , $j = 1, 2, 3, 4, 5$, and their partial derivatives. After eliminating any dependencies among the derivatives of the m 's caused by the system itself, we can then equate the coefficients of the remaining unconstrained partial derivatives of u to zero. This will result

in a large number of elementary partial differential equations for the coefficient functions ξ_i , η_j of the infinitesimal generator, called the determining equations for the symmetry group of the given system. In this case, we find the determining equations for the symmetry group of the system (1) to be the following:

$$\begin{aligned}
&\xi_{1,t^2} = \xi_{1,x} = \xi_{1,y} = \xi_{1,z} = \xi_{1,u} = \xi_{1,v} = \xi_{1,w} = \xi_{1,q} = \xi_{1,p} = 0, \\
&\xi_{2,x} = \xi_{2,u} = \xi_{2,v} = \xi_{2,w} = \xi_{2,q} = \xi_{2,p} = \xi_{2,t^2} = \xi_{2,ty} = \xi_{2,y^2} = 0, \\
&\xi_{3,y} = \xi_{3,u} = \xi_{3,v} = \xi_{3,w} = \xi_{3,q} = \xi_{3,p} = \xi_{3,t^2} = 0, \\
&\xi_{4,z} = \xi_{4,w} = \xi_{4,q} = \xi_{4,p} = 0, \\
&\xi_{4,t^2} = \xi_{4,tx} = \xi_{4,ty} = \xi_{4,x^2} = \xi_{4,xy} = \xi_{4,y^2} = 0, \\
&\xi_{3,x} = \xi_{2,y}, \xi_{3,z} + \xi_{4,y} = 0, \xi_{4,v} = \xi_{4,u}, \xi_{2,z} + \xi_{4,x} = 0,
\end{aligned} \tag{3}$$

and

$$\begin{aligned}
&\eta_1 = -u\xi_{1,t} + \xi_{2,t} + v\xi_{2,y} - w\xi_{4,x}, \\
&\eta_2 = -w\xi_{4,y} + \xi_{3,t} - u\xi_{2,y} - v\xi_{1,t}, \\
&\eta_3 = u\xi_{4,x} + \xi_{4,t} + v\xi_{4,y} - w\xi_{1,t}, \\
&\eta_4 = q\eta_5/p + 2q\xi_{1,t}, \\
&\eta_{5,t} = \eta_{5,x} = \eta_{5,y} = \eta_{5,z} = \eta_{5,u} = \eta_{5,v} = \eta_{5,w} = \eta_{5,q} = 0, \\
&\eta_5 = p\eta_{5,p}.
\end{aligned} \tag{4}$$

First, equations (3) require that ξ_i s are just

$$\begin{aligned}
&\xi_1 = a_1 t + a_2, \\
&\xi_2 = a_3 t + a_4 x + a_5 y + a_6 z + a_7, \\
&\xi_3 = a_8 t - a_5 x + a_4 y + a_9 z + a_{10}, \\
&\xi_4 = a_{11} t - a_6 x - a_9 y + a_4 z + a_{12},
\end{aligned} \tag{5}$$

where $a_1, \dots, a_{12}$ are arbitrary constants. Then, equations (4) and (5) requires that

$$\begin{aligned}
&\eta_1 = (a_4 - a_1)u + a_5 v - a_6 w + a_3, \\
&\eta_2 = -a_5 u + (a_4 - a_1)v + a_4 w + a_8, \\
&\eta_3 = -a_6 u - a_9 v + (a_4 - a_1)w + a_{11}, \\
&\eta_4 = (2a_1 - 2a_4 + a_{13})q, \\
&\eta_5 = a_{13}p,
\end{aligned} \tag{6}$$

where a_{13} is an arbitrary constant. Therefore

Theorem 1.. The Lie algebra of infinitesimal symmetries of the system (1) is spanned by the 13 vector fields

$$\begin{aligned}
& X_1 = \partial_x, \\
& X_2 = \partial_y, \\
& X_3 = \partial_z, \\
& X_4 = \partial_t, \\
& \hspace{15em} (\text{shifts of origin}) \\
& X_5 = t \partial_x + \partial_u, \\
& X_6 = t \partial_y + \partial_v, \\
& X_7 = t \partial_z + \partial_w, \\
& \hspace{15em} (\text{uniformly frame motions}) \\
& X_8 = t \partial_t - u \partial_u - v \partial_v - w \partial_w + 2q \partial_q, \\
& X_9 = x \partial_x + y \partial_y + z \partial_z + u \partial_u + v \partial_v + w \partial_w - 2r \partial_r, \\
& X_{10} = y \partial_x - x \partial_y + v \partial_u - u \partial_v, \\
& X_{11} = -z \partial_y + y \partial_z - w \partial_v + v \partial_w, \\
& X_{12} = -z \partial_x + x \partial_z - w \partial_u + u \partial_w, \\
& \hspace{15em} (\text{rotations of reference frame}) \\
& X_{13} = q \partial_q + p \partial_p,
\end{aligned} \tag{7}$$

These infinitesimal symmetries will generate a Lie algebra $\mathfrak{g}$ over the field of real or complex numbers. The commutator table of Lie algebra $\mathfrak{g}$ for (1) is given below, where the entry in the i th row and j th column is defined as $[X_i, X_j] = X_i.X_j - X_j.X_i$, $i, j = 1, \dots, 13$.

	X_1	X_2	X_3	X_4	X_5	X_6	X_7	X_8	X_9	X_{10}	X_{11}	X_{12}	X_{13}
X_1	0	0	0	0	0	0	0	0	X_1	$-X_2$	0	X_3	0
X_2	0	0	0	0	0	0	0	0	X_2	X_1	X_3	0	0
X_3	0	0	0	0	0	0	0	0	X_3	0	$-X_2$	$-X_1$	0
X_4	0	0	0	0	X_1	X_2	X_3	X_4	0	0	0	0	0
X_5	0	0	0	$-X_1$	0	0	0	$-X_5$	X_5	$-X_6$	0	X_7	0
X_6	0	0	0	$-X_2$	0	0	0	$-X_6$	X_6	X_5	X_7	0	0
X_7	0	0	0	$-X_3$	0	0	0	$-X_7$	X_7	0	$-X_6$	$-X_5$	0
X_8	0	0	0	$-X_4$	X_5	X_6	X_7	0	0	0	0	0	0
X_9	$-X_1$	$-X_2$	$-X_3$	0	$-X_5$	$-X_6$	$-X_7$	0	0	0	0	0	0
X_{10}	X_2	$-X_1$	0	0	X_6	$-X_5$	0	0	0	0	$-X_{12}$	X_{11}	0
X_{11}	0	$-X_3$	X_2	0	0	$-X_7$	X_6	0	0	X_{12}	0	$-X_{10}$	0
X_{12}	$-X_3$	0	X_1	0	$-X_7$	0	X_5	0	0	$-X_{11}$	X_{10}	0	0
X_{13}	0	0	0	0	0	0	0	0	0	0	0	0	0

3. Reduction of the system

Now, we look at the group-invariant solutions to the system (1).

The system (1) is a sub-manifold M of jet space $J^2(\mathbb{R}^4, \mathbb{R}^5)$ defined by the following equations

$$\begin{aligned}
u_t + uu_x + vu_y + wu_z + p_x/q &= 0, \\
v_t + uv_x + vv_y + wv_z + p_y/q &= 0, \\
w_t + uw_x + vw_y + ww_z + p_z/q &= 0, \\
qt + q(u_x + v_y + w_z) + uq_x + vq_y + wq_z &= 0, \\
p_t + \gamma p(u_x + v_y + w_z) + up_x + vp_y + wp_z &= 0.
\end{aligned} \tag{8}$$

Doing as section 3.1 of [6] and find (in a sense) the most general group-invariant solutions to the system (1).

Theorem 2.. The one-parameter groups $g_i(s) : M \rightarrow M$ generated by the X_i , $i = 1, \dots, 13$ are given in the following table:

$$\begin{aligned}
g_1 &: (t, x + s, y, z, u, v, w, p, q) \\
g_2 &: (t, x, y + s, z, u, v, w, p, q) \\
g_3 &: (t, x, y, z + s, u, v, w, p, q) \\
g_4 &: (t + s, x, y, z, u, v, w, p, q) \\
g_5 &: (t, x + st, y, z, u + s, v, w, p, q) \\
g_6 &: (t, x, y + st, z, u, v + s, w, p, q) \\
g_7 &: (t, x, y, z + st, u, v, w + s, p, q) \\
g_8 &: (e^s t, x, y, z, e^{-s} u, e^{-s} v, e^{-s} w, p, e^{2s} q) \\
g_9 &: (t, e^s x, e^s y, e^s z, e^s u, e^s v, e^s w, p, e^{-2s} q) \\
g_{10} &: (t, x \cos s + y \sin s, y \cos s - x \sin s, z, \\
&\quad v \sin s + u \cos s, v \cos s - u \sin s, w, p, q) \\
g_{11} &: (t, x, y \cos s - z \sin s, z \cos s + y \sin s, \\
&\quad u, v \cos s - w \sin s, w \cos s + v \sin s, p, q) \\
g_{12} &: (t, x \cos s - z \sin s, y, x \sin s + z \cos s, \\
&\quad u \cos s - w \sin s, v, w \cos s + u \sin s, p, q) \\
g_{13} &: (t, x, y, z, u, v, w, e^s p, e^s q)
\end{aligned} \tag{9}$$

Where, entries give the transformed point

$$\exp(sX_i)(t, x, y, z, u, v, w, p, q) = (\tilde{t}, \tilde{x}, \tilde{y}, \tilde{z}, \tilde{u}, \tilde{v}, \tilde{w}, \tilde{p}, \tilde{q}).$$

Since each group g_i , $i = 1, \dots, 13$, is a symmetry group of (1), then

Theorem 3.. Let $i = 1, \dots, 13$, $s \in \mathbb{R}$, and $\lambda \in \mathbb{R} - \{0\}$. If

$$\begin{aligned} u &= U(t, x, y, z), \quad v = V(t, x, y, z), \quad w = W(t, x, y, z), \\ p &= P(t, x, y, z), \quad q = Q(t, x, y, z), \end{aligned}$$

is a solution of the system (1), so are the functions

$$\begin{aligned} u^{(i)} &= U^{(i)}(t, x, y, z), \quad v^{(i)} = V^{(i)}(t, x, y, z), \quad w^{(i)} = W^{(i)}(t, x, y, z), \\ p^{(i)} &= P^{(i)}(t, x, y, z), \quad q^{(i)} = Q^{(i)}(t, x, y, z), \end{aligned}$$

where

$$\begin{aligned} u^{(1)} &= U(t, x + s, y, z), & u^{(2)} &= U(t, x, y + s, z), \\ v^{(1)} &= V(t, x + s, y, z), & v^{(2)} &= V(t, x, y + s, z), \\ w^{(1)} &= W(t, x + s, y, z), & w^{(2)} &= W(t, x, y + s, z), \\ p^{(1)} &= P(t, x + s, y, z), & p^{(2)} &= P(t, x, y + s, z), \\ q^{(1)} &= Q(t, x + s, y, z). & q^{(2)} &= Q(t, x, y + s, z). \end{aligned}$$

$$\begin{aligned} u^{(3)} &= U(t, x, y, z + s), & u^{(4)} &= U(t + s, x, y, z), \\ v^{(3)} &= V(t, x, y, z + s), & v^{(4)} &= V(t + s, x, y, z), \\ w^{(3)} &= W(t, x, y, z + s), & w^{(4)} &= W(t + s, x, y, z), \\ p^{(3)} &= P(t, x, y, z + s), & p^{(4)} &= P(t + s, x, y, z), \\ q^{(3)} &= Q(t, x, y, z + s). & q^{(4)} &= Q(t + s, x, y, z). \end{aligned}$$

$$\begin{aligned} u^{(5)} &= U(t, x + st, y, z) - s, & u^{(6)} &= U(t, x, y + st, z), \\ v^{(5)} &= V(t, x + st, y, z), & v^{(6)} &= V(t, x, y + st, z) - s, \\ w^{(5)} &= W(t, x + st, y, z), & w^{(6)} &= W(t, x, y + st, z), \\ p^{(5)} &= P(t, x + st, y, z), & p^{(6)} &= P(t, x, y + st, z), \\ q^{(5)} &= Q(t, x + st, y, z). & q^{(6)} &= Q(t, x, y + st, z). \end{aligned}$$

$$\begin{aligned} u^{(7)} &= U(t, x, y, z + st), & u^{(8)} &= \lambda U(\lambda.t, x, y, z), \\ v^{(7)} &= V(t, x, y, z + st), & v^{(8)} &= \lambda V(\lambda.t, x, y, z), \\ w^{(7)} &= W(t, x, y, z + st) - s, & w^{(8)} &= \lambda W(\lambda.t, x, y, z), \\ p^{(7)} &= P(t, x, y, z + st), & p^{(8)} &= P(\lambda.t, x, y, z), \\ q^{(7)} &= Q(t, x, y, z + st). & q^{(8)} &= \frac{1}{\lambda^2} Q(\lambda.t, x, y, z). \end{aligned}$$

$$\begin{aligned} u^{(9)} &= \frac{1}{\lambda} U(t, \lambda.x, \lambda.y, \lambda.z), & u^{(13)} &= U(t, x, y, z), \\ v^{(9)} &= \frac{1}{\lambda} V(t, \lambda.x, \lambda.y, \lambda.z), & v^{(13)} &= V(t, x, y, z), \\ w^{(9)} &= \frac{1}{\lambda} W(t, \lambda.x, \lambda.y, \lambda.z), & w^{(13)} &= W(t, x, y, z), \\ p^{(9)} &= P(t, \lambda.x, \lambda.y, \lambda.z), & p^{(13)} &= \lambda P(t, x, y, z), \\ q^{(9)} &= \lambda^2 Q(t, \lambda.x, \lambda.y, \lambda.z). & q^{(13)} &= \lambda Q(t, x, y, z). \end{aligned}$$

$$\begin{aligned}
u^{(10)} &= \cos s.U(t, x \cos s + y \sin s, y \cos s - x \sin s, z) \\
&\quad - \sin s.V(t, x \cos s + y \sin s, y \cos s - x \sin s, z), \\
v^{(10)} &= \sin s.U(t, x \cos s + y \sin s, y \cos s - x \sin s, z) \\
&\quad + \cos s.V(t, x \cos s + y \sin s, y \cos s - x \sin s, z), \\
w^{(10)} &= W(t, x \cos s + y \sin s, y \cos s - x \sin s, z), \\
p^{(10)} &= P(t, x \cos s + y \sin s, y \cos s - x \sin s, z), \\
q^{(10)} &= Q(t, x \cos s + y \sin s, y \cos s - x \sin s, z).
\end{aligned}$$

$$\begin{aligned}
u^{(11)} &= U(t, x, y \cos s - z \sin s, z \cos s + y \sin s), \\
v^{(11)} &= \cos s.V(t, x, y \cos s - z \sin s, z \cos s + y \sin s) \\
&\quad + \sin s.W(t, x, y \cos s - z \sin s, z \cos s + y \sin s), \\
w^{(11)} &= \cos s.W(t, x, y \cos s - z \sin s, z \cos s + y \sin s) \\
&\quad - \sin s.V(t, x, y \cos s - z \sin s, z \cos s + y \sin s), \\
p^{(11)} &= P(t, x, y \cos s - z \sin s, z \cos s + y \sin s), \\
q^{(11)} &= Q(t, x, y \cos s - z \sin s, z \cos s + y \sin s).
\end{aligned}$$

$$\begin{aligned}
u^{(12)} &= \cos s.U(t, x \cos s - z \sin s, y, x \sin s + z \cos s) \\
&\quad + \sin s.W(t, x \cos s - z \sin s, y, x \sin s + z \cos s), \\
v^{(12)} &= V(t, x \cos s - z \sin s, y, x \sin s + z \cos s), \\
w^{(12)} &= \cos s.W(t, x \cos s - z \sin s, y, x \sin s + z \cos s) \\
&\quad - \sin s.U(t, x \cos s - z \sin s, y, x \sin s + z \cos s), \\
p^{(12)} &= P(t, x \cos s - z \sin s, y, x \sin s + z \cos s), \\
q^{(12)} &= Q(t, x \cos s - z \sin s, y, x \sin s + z \cos s).
\end{aligned}$$

4. Structure of Lie algebra $\mathfrak{g}$

In this section, we determine the structure of full symmetry algebra $\mathfrak{g}$ of system (1).

The center $\mathfrak{z}$ of $\mathfrak{g}$ is $\text{Span}_{\mathbb{R}}\{X_{13}\}$. Therefore, the quotient algebra $\mathfrak{g}_1 := \mathfrak{g}/\mathfrak{z}$ is $\text{Span}_{\mathbb{R}}\{Y_1, \dots, Y_{12}\}$; where $Y_i := X_i + \mathfrak{z}$, $i = 1, \dots, 12$. The commutator table of Lie algebra $\mathfrak{g}_1$ is given below, where the entry in the i th row and j th column

is defined as $[Y_i, Y_j] = Y_i \cdot Y_j - Y_j \cdot Y_i$, $i, j = 1, \dots, 12$.

	Y_1	Y_2	Y_3	Y_4	Y_5	Y_6	Y_7	Y_8	Y_9	Y_{10}	Y_{11}	Y_{12}
Y_1	0	0	0	0	0	0	0	0	Y_1	$-Y_2$	0	Y_3
Y_2	0	0	0	0	0	0	0	0	Y_2	Y_1	Y_3	0
Y_3	0	0	0	0	0	0	0	0	Y_3	0	$-Y_2$	$-Y_1$
Y_4	0	0	0	0	Y_1	Y_2	Y_3	Y_4	0	0	0	0
Y_5	0	0	0	$-Y_1$	0	0	0	$-Y_5$	Y_5	$-Y_6$	0	Y_7
Y_6	0	0	0	$-Y_2$	0	0	0	$-Y_6$	Y_6	Y_5	Y_7	0
Y_7	0	0	0	$-Y_3$	0	0	0	$-Y_7$	Y_7	0	$-Y_6$	$-Y_5$
Y_8	0	0	0	$-Y_4$	Y_5	Y_6	Y_7	0	0	0	0	0
Y_9	$-Y_1$	$-Y_2$	$-Y_3$	0	$-Y_5$	$-Y_6$	$-Y_7$	0	0	0	0	0
Y_{10}	Y_2	$-Y_1$	0	0	Y_6	$-Y_5$	0	0	0	0	$-Y_{12}$	Y_{11}
Y_{11}	0	$-Y_3$	Y_2	0	0	$-Y_7$	Y_6	0	0	Y_{12}	0	$-Y_{10}$
Y_{12}	$-Y_3$	0	Y_1	0	$-Y_7$	0	Y_5	0	0	$-Y_{11}$	Y_{10}	0

The algebra $\mathfrak{g}_1$ is not solvable, because

$$\mathfrak{g}_1^{(1)} := [\mathfrak{g}_1, \mathfrak{g}_1] = \text{Span}_{\mathbb{R}}\{Y_1, \dots, Y_{10}\}, \text{ and } \mathfrak{g}_1^{(2)} := [\mathfrak{g}_1^{(1)}, \mathfrak{g}_1^{(1)}] = \mathfrak{g}_1^{(1)}.$$

Lie algebra $\mathfrak{g}_1$ admits a Levi-decomposition into the semi-direct sum

$$\mathfrak{g}_1 = \mathfrak{r} \ltimes \mathfrak{s},$$

where $\mathfrak{r} = \text{Span}_{\mathbb{R}}\{Y_1, \dots, Y_9\}$ is the radical of $\mathfrak{g}_1$ (the largest solvable ideal contained in $\mathfrak{g}_1$) and $\mathfrak{s} = \text{Span}_{\mathbb{R}}\{Y_{10}, Y_{11}, Y_{12}\}$ is a semi-simple subalgebra of $\mathfrak{g}_1$. The radical $\mathfrak{r}$ is uniquely defined but the semi-simple subalgebra $\mathfrak{s}$ is not. $\mathfrak{s}$ as an algebra is isomorphic to 3-dimensional special orthogonal algebra $\mathfrak{so}(3) := \{A \in \text{Mat}(3 \times 3; \mathbb{R}) \mid A^t = -A\}$, which is a simple algebra.

The radical $\mathfrak{r}$ is solvable, with and

$$\mathfrak{r}^{(1)} = \text{Span}_{\mathbb{R}}\{Y_1, \dots, Y_7\}, \quad \mathfrak{r}^{(2)} = \text{Span}_{\mathbb{R}}\{Y_1, Y_2, Y_3\}, \quad \mathfrak{r}^{(3)} = \{0\}.$$

It is semi-direct sum $\mathfrak{r} = \mathfrak{r}^{(1)} \ltimes \mathfrak{g}_2$, where $\mathfrak{g}_2 = \text{Span}_{\mathbb{R}}\{Y_8, Y_9\}$ is isomorphic to the two dimensional Abelian Lie algebra $\mathbb{R}^2$; i.e.

$$\mathfrak{r} = \mathfrak{r}^{(1)} \ltimes \mathbb{R}^2.$$

$\mathfrak{r}^{(1)}$ is semi-direct sum $\mathfrak{r}^{(1)} = \mathfrak{r}^{(2)} \ltimes \mathfrak{g}_3$, where $\mathfrak{g}_3 = \text{Span}_{\mathbb{R}}\{Y_4, \dots, Y_7\}$ is isomorphic to the four dimensional Abelian Lie algebra $\mathbb{R}^4$, and $\mathfrak{r}^{(2)}$ is isomorphic to the three dimensional Abelian Lie algebra $\mathbb{R}^3$; i.e.

$$\mathfrak{r}^{(1)} = \mathbb{R}^3 \ltimes \mathbb{R}^4.$$

Theorem 4.. The symmetry algebra $\mathfrak{g}$ of system (1) have the following structure

$$\mathfrak{g} \cong \mathbb{R} \times ((\mathbb{R}^3 \ltimes \mathbb{R}^4) \ltimes \mathbb{R}^2) \ltimes \mathfrak{so}(3).$$

Theorem 5.. There is a normal Lie-subalgebra of symmetry Lie-group of system (1) which is isomorphic to the Galilean group of $\mathbb{R}^4$:

$$\text{Gal}(4, \mathbb{R}) = \left\{ \begin{pmatrix} 1 & \mathbf{0} & s \\ \mathbf{v} & \mathbf{R} & \mathbf{y} \\ 0 & \mathbf{0} & 1 \end{pmatrix} \mid \mathbf{R} \in \text{O}(3, \mathbb{R}), s \in \mathbb{R}, \text{ and } \mathbf{y}, \mathbf{v} \in \mathbb{R}^3 \right\}.$$

Therefore, the system (1) is invariant up to Galilean motions of space-times $\mathbb{R}^4$ (see [3]).

Proof. Let $\mathfrak{b} = \text{Span}_{\mathbb{R}}\{X_1, \dots, X_7, X_{10}, X_{11}, X_{12}\}$. Lie algebra structure of $\mathfrak{b}$ is as $\mathfrak{gal}(3)$, the lie algebra of Galilean group $\text{Gal}(4, \mathbb{R})$, and $\mathfrak{b}$ an ideal of $\mathfrak{g}$. Therefore, there is a Lie-subgroup of G such that its Lie-algebra is $\mathfrak{b}$, by the Theorem 2.53 of Olver [6]. $\square$

Conclusion 1.. $\mathfrak{g}$ is semi-direct sum of

$$\mathfrak{gal}(4) \cong \text{Span}_{\mathbb{R}}\{X_1, \dots, X_7, X_{10}, X_{11}, X_{12}\}$$

and 3-dimensional Abelian algebra $\mathbb{R}^3 \cong \text{Span}\{X_8, X_9, X_{13}\}$.

5. Optimal system of sub-algebras

As is well known, the Lie group theoretic method plays an important role in finding exact solutions and performing symmetry reductions of differential equations. Since any linear combination of infinitesimal generators is also an infinitesimal generator, there are always infinitely many different symmetry subgroups for the differential equation. So, a mean of determining which subgroups would give essentially different types of solutions is necessary and significant for a complete understanding of the invariant solutions. As any transformation in the full symmetry group maps a solution to another solution, it is sufficient to find invariant solutions which are not related by transformations in the full symmetry group, this has led to the concept of an optimal system [7]. The problem of finding an optimal system of subgroups is equivalent to that of finding an optimal system of subalgebras. For one-dimensional subalgebras, this classification problem is essentially the same as the problem of classifying the orbits of the adjoint representation. This problem is attacked by the naive approach of taking a general element in the Lie algebra and subjecting it to various adjoint transformations so as to simplify it as much as possible. The idea of using the adjoint representation to classify group-invariant solutions was due to [7] and [5].

The adjoint action is given by the Lie series

$$\text{Ad}(\exp(sX_i)X_j) = X_j - s[X_i, X_j] + \frac{s^2}{2}[X_i, [X_i, X_j]] - \dots, \quad (10)$$

where $[X_i, X_j]$ is the commutator for the Lie algebra, s is a parameter, and $i, j = 1, \dots, 13$. We can write the adjoint action for the Lie algebra $\mathfrak{g}$, and show that

Theorem 6.. A one-dimensional optimal system of (1) is given by

- 1) $X = X_8 + \sum_{i=9}^{13} a_i X_i,$
- 2) $X = a_7 X_7 + X_8 + X_9 + \sum_{i=10}^{13} a_i X_i,$
- 3) $X = a_6 X_6 + a_7 X_7 + X_8 + X_9 + \sum_{i=11}^{13} a_i X_i,$
- 4) $X = a_5 X_5 + X_8 + X_9 + a_{11} X_i + a_{13} X_{13},$
- 5) $X = \sum_{i=5}^7 a_i X_i + X_8 + X_9 + a_{13} X_{13},$
- 6) $X = a_4 X_4 + X_9 + \sum_{i=10}^{13} a_i X_i,$
- 7) $X = X_4 + a_5 X_5 + a_8 X_8 + \sum_{i=10}^{13} a_i X_i,$
- 8) $X = X_4 + a_6 X_6 + a_8 X_8 + a_{10} X_{10} + a_{12} X_{12} + a_{13} X_{13},$
- 9) $X = X_4 + a_7 X_7 + a_8 X_8 + a_{10} X_{10} + a_{12} X_{12} + a_{13} X_{13},$
- 10) $X = X_4 + \sum_{i=5}^8 a_i X_i + a_{13} X_{13},$
- 11) $X = X_4 + a_7 X_7 + \sum_{i=10}^{13} a_i X_i,$
- 12) $X = X_7 + \sum_{i=10}^{13} a_i X_i,$
- 13) $X = a_3 X_3 + \sum_{i=10}^{13} a_i X_i,$
- 14) $X = X_4 + a_6 X_6 + \sum_{i=11}^{13} a_i X_i,$
- 15) $X = X_4 + a_5 X_5 + a_{11} X_{11} + a_{13} X_{13},$
- 16) $X = X_4 + \sum_{i=5}^7 a_i X_i + a_{13} X_{13},$
- 17) $X = X_6 + \sum_{i=11}^{13} a_i X_i,$
- 18) $X = a_2 X_2 + \sum_{i=11}^{13} a_i X_i,$
- 19) $X = a_2 X_2 + a_{12} X_{12} + a_{13} X_{13},$
- 20) $X = X_5 + a_{11} X_{11} + a_{13} X_{13},$
- 21) $X = a_2 X_2 + a_3 X_3 + X_5 + a_{13} X_{13},$
- 22) $X = a_1 X_1 + X_{11} + a_{13} X_{13},$
- 23) $X = a_1 X_1 + a_3 X_3 + X_6 + a_{13} X_{13},$
- 24) $X = a_1 X_1 + a_2 X_2 + a_3 X_3 + a_{13} X_{13},$
- 25) $X = X_5 + a_{10} X_{10} + a_{11} X_{11} + a_{13} X_{13},$

$$26) \quad X = a_3 X_3 + X_{10} + a_{11} X_{11} + a_{13} X_{13}.$$

Proof: $F(sX_i) : \mathfrak{g} \rightarrow \mathfrak{g}$ defined by $X \mapsto \text{Ad}(\exp(sX_i)X)$ is a linear map, for $i = 1, \dots, 13$. The matrix M_{sX_i} of $F(sX_i)$, $i = 1, \dots, 13$, with respect to basis $\{X_1, \dots, X_{13}\}$ is

[illegible]

[illegible]

[illegible]

[illegible]

[illegible]

[illegible]

[illegible]

12

$$\begin{bmatrix}
1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & \cos s & \sin s & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & -\sin s & \cos s & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & \cos s & \sin s & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & -\sin s & \cos s & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & \cos s & 0 & -\sin s & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & \sin s & 0 & \cos s & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1
\end{bmatrix},$$

$$\begin{bmatrix}
\cos s & 0 & \sin s & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\
-\sin s & 0 & \cos s & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & \cos s & 0 & \sin s & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & -\sin s & 0 & \cos s & 0 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & \cos s & \sin s & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & -\sin s & \cos s & 0 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\
0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1
\end{bmatrix},$$

and I_{13} , respectively. Let $X = \sum_{i=1}^{13} a_i X_i$, then

$$\begin{aligned}
& F(s_7.X_7) \circ F(s_6.X_6) \circ \cdots \circ F(s_1.X_1) : X \mapsto \\
& (s_3 a_{12} - s_2 a_{10} - s_1 a_9 + a_1 - s_5 s_4 a_8 + s_5 a_4 - s_4 a_5).X_1 \\
& + (-s_6 s_4 a_8 + s_3 a_{11} - s_2 a_9 + s_1 a_{10} + a_2 + s_6 a_4 - s_4 a_6).X_2 \\
& + (-s_7 s_4 a_8 - s_3 a_9 - s_2 a_{11} - s_1 a_{12} + a_3 + s_7 a_4 - s_4 a_7).X_3 \\
& + (a_4 - s_4 a_8).X_4 + (-s_6 a_{10} - s_5 a_9 + s_5 a_8 + a_5 + s_7 a_{12}).X_5 \\
& + (-s_6 a_9 + s_6 a_8 + s_5 a_{10} + a_6 + s_7 a_{11}).X_6 \\
& + (s_7 a_8 - s_6 a_{11} - s_5 a_{12} + a_7 - s_7 a_9).X_7 \\
& + a_8.X_8 + a_9.X_9 + a_{10}.X_{10} + a_{11}.X_{11} + a_{12}.X_{12} + a_{13}.X_{13}
\end{aligned} \tag{11}$$

If a_8 , and $a_9 \neq 0$, and $a_8 \neq 2a_9$, then we can make the coefficients of $X_1, \dots, X_7$ vanish. Scaling X if necessary, we can assume that $a_8 = 1$. And X is reduced to Case 1.

If $a_8 = a_9 \neq 0$, and $a_{10} \neq 0$, then we can make the coefficients of $X_1, \dots, X_6$ vanish. Scaling X if necessary, we can assume that $a_8 = 1$. And X is reduced to Case 2.

If $a_8 = a_9 \neq 0$, $a_{10} = 0$, and $a_{12} \neq 0$, then we can make the coefficients of $X_1, \dots, X_5$, and X_{10} vanish. Scaling X if necessary, we can assume that $a_8 = 1$. And X is reduced to Case 3.

If $a_8 = a_9 \neq 0$, $a_{10} = a_{12} = 0$, and $a_{11} \neq 0$, then we can make the coefficients of $X_1, \dots, X_4$, X_6 , X_7 , X_{10} , and X_{12} vanish. Scaling X if necessary, we can assume that $a_8 = 1$. And X is reduced to Case 4.

If $a_8 = a_9 \neq 0$, $a_{10} = a_{11} = a_{12} = 0$, then we can make the coefficients of $X_1, \dots, X_4$, and $X_{10}, \dots, X_{11}$ vanish. Scaling X if necessary, we can assume that $a_8 = 1$. And X is reduced to Case 5.

If $a_8 = 0$, and $a_9 \neq 0$, then we can make the coefficients of $X_1, \dots, X_3$, and $X_5, \dots, X_8$ vanish. Scaling X if necessary, we can assume that $a_9 = 1$. And X

is reduced to Case 6.

If $a_8 \neq 0$, $a_9 = 0$, and $a_{11} \neq 0$, then we can make the coefficients of $X_1, \dots, X_3$, and X_6, X_7 , and X_9 vanish. And X is reduced to Case 7.

If $a_8 \neq 0$, $a_9 = a_{11} = 0$, and $a_{12} \neq 0$, then we can make the coefficients of $X_1, \dots, X_3$, and X_5, X_7, X_9 , and X_{11} vanish. And X is reduced to Case 8.

If $a_8 \neq 0$, $a_9 = a_{11} = a_{12} = 0$, and $a_{10} \neq 0$, then we can make the coefficients of $X_1, \dots, X_3$, and X_5, X_6, X_9 , and X_{11} vanish. And X is reduced to Case 9.

If $a_8 \neq 0$, and $a_9 = a_{10} = a_{11} = a_{12} = 0$, then we can make the coefficients of $X_1, \dots, X_3$, and $X_9, \dots, X_{12}$ vanish. And X is reduced to Case 10.

If $a_8 = a_9 = 0$, $a_4 \neq 0$, and $a_{10} \neq 0$, then we can make the coefficients of $X_1, \dots, X_3, X_5, X_6, X_8$, and X_9 vanish. Scaling X if necessary, we can assume that $a_4 = 1$. And X is reduced to Case 11.

If $a_4 = a_8 = a_9 = 0$, $a_{10} \neq 0$, $a_{12} \neq 0$, and $a_6 a_{12} + a_7 a_{10} \neq a_5 a_{11}$, then we can make the coefficients of $X_1, \dots, X_6, X_8$, and X_9 vanish. Scaling X if necessary, we can assume that $a_7 = 1$. And X is reduced to Case 12.

If $a_4 = a_8 = a_9 = 0$, $a_{10} \neq 0$, $a_{12} \neq 0$, and $a_6 a_{12} + a_7 a_{10} = a_5 a_{11}$, then we can make the coefficients of X_1, X_2 , and $X_4, \dots, X_9$ vanish. And X is reduced to Case 13.

If $a_8 = a_9 = a_{10} = 0$, and $a_4 \neq 0$, then we can make the coefficients of $X_1, \dots, X_3, X_5$, and $X_7, \dots, X_{10}$ vanish. Scaling X if necessary, we can assume that $a_4 = 1$. And X is reduced to Case 14.

If $a_8 = \dots = a_{10} = 0$, $a_{12} = 0$, $a_4 \neq 0$, and $a_{11} \neq 0$, then we can make the coefficients of $X_1, \dots, X_3$, and $X_8, \dots, X_{10}$ vanish. Scaling X if necessary, we can assume that $a_4 = 1$. And X is reduced to Case 15.

If $a_8 = \dots = a_{12} = 0$, and $a_4 \neq 0$, then we can make the coefficients of $X_1, \dots, X_3$, and $X_8, \dots, X_{10}$ vanish. Scaling X if necessary, we can assume that $a_4 = 1$. And X is reduced to Case 16.

If $a_4 = a_8 = a_9 = a_{10} = 0$, $a_{12} \neq 0$, and $a_5 a_{11} \neq a_6 a_{12}$, then we can make the coefficients of $X_1, \dots, X_5$, and $X_7, \dots, X_{10}$ vanish. Scaling X if necessary, we can assume that $a_6 = 1$. And X is reduced to Case 17.

If $a_4 = a_8 = a_9 = a_{10} = 0$, $a_{12} \neq 0$, $a_5 a_{11} = a_6 a_{12}$, and $a_6 \neq 0$, then we can make the coefficients of $X_1, \dots, X_5$, and $X_7, \dots, X_{10}$ vanish. And X is reduced to Case 18.

If $a_4 = a_6 = a_8 = a_9 = a_{10} = 0$, $a_{12} \neq 0$, $a_5 a_{11} = a_6 a_{12}$, and $a_6 \neq 0$, then we can make the coefficients of $X_1, \dots, X_5$, and $X_7, \dots, X_{10}$ vanish. And X is reduced to Case 18.

If $a_4 = a_6 = a_8 = \dots = a_{11} = 0$, $a_{12} \neq 0$, $a_5 a_{11} = a_6 a_{12}$, $a_6 \neq 0$, and $a_5 \neq 0$, then we can make the coefficients of X_1 , and $X_3, \dots, X_{11}$ vanish. And X is reduced to Case 19.

If $a_4 = a_8 = a_9 = a_{10} = a_{12} = 0$, $a_5 \neq 0$, and $a_{11} \neq 0$, then we can make the coefficients of $X_1, \dots, a_4, X_6, \dots, X_{10}$, and a_{12} vanish. Scaling X if necessary, we can assume that $a_5 = 1$. And X is reduced to Case 20.

If $a_4 = a_8 = \dots = a_{12} = 0$, and $a_5 \neq 0$, then we can make the coefficients of X_1, a_4 , and $X_8, \dots, X_{12}$ vanish. Scaling X if necessary, we can assume that

$a_5 = 1$. And X is reduced to Case 21.

If $a_4 = a_5 = a_8 = a_9 = a_{10} = a_{12} = 0$, and $a_{11} \neq 0$, then we can make the coefficients of $X_2, \dots, X_{10}$ and a_{12} vanish. Scaling X if necessary, we can assume that $a_{11} = 1$. And X is reduced to Case 22.

If $a_4 = a_5 = a_8 = \dots = a_{12} = 0$, and $a_6 \neq 0$, then we can make the coefficients of X_2, a_5 , and $X_7, \dots, X_{12}$ vanish. Scaling X if necessary, we can assume that $a_6 = 1$. And X is reduced to Case 23.

If $a_4 = a_5 = a_6 = 0$, and $a_8 = \dots = a_{12} = 0$, then we can make the coefficients of $X_4, \dots, X_{12}$ vanish. And X is reduced to Case 24.

If $a_4 = a_8 = a_9 = a_{12} = 0$, $a_{10} \neq 0$, and $a_7 a_{10} \neq a_5 a_{11}$, then we can make the coefficients of $X_1, \dots, X_4, X_6, \dots, X_9$, and $a_{12} = 0$ vanish. Scaling X if necessary, we can assume that $a_5 = 1$. And X is reduced to Case 25.

If $a_4 = a_8 = a_9 = a_{12} = 0$, $a_7 a_{10} = a_5 a_{11}$, and $a_{10} \neq 0$, then we can make the coefficients of $X_1, \dots, X_4, X_6, \dots, X_9$, and $a_{12} = 0$ vanish. Scaling X if necessary, we can assume that $a_{10} = 1$. And X is reduced to Case 26. And the Theorem follows. $\square$

According to our optimal system of one-dimensional subalgebras of the full symmetry algebra $\mathfrak{g}$, we need only find group-invariant solutions for 26 one-parameter subgroups generated by X as Theorem 6. For example, as a direct consequence of Theorem 3 and case 1 of Theorem 6, we deduce the following conclusion:

Conclusion 2. Let $s, c_1, \dots, c_6 \in \mathbb{R}$ with c_1, c_2 , and $c_6 \neq 0$, and s be sufficiently small. Then, if

$$\begin{aligned} u &= U(t, x, y, z), \quad v = V(t, x, y, z), \quad w = W(t, x, y, z), \\ p &= P(t, x, y, z), \quad q = Q(t, x, y, z), \end{aligned}$$

is a solution of the system (1), so are the functions

$$\begin{aligned} \tilde{u} &= c_1^s c_2^{-s} (\cos(c_3 s) \cos(c_5 s) - \sin(c_3 s) \sin(c_4 s) \sin(c_5 s)) \cdot U(\tilde{t}, \tilde{x}, \tilde{y}, \tilde{z}) \\ &\quad - c_1^s c_2^{-s} (\sin(c_3 s) \cos(c_5 s) + \cos(c_3 s) \sin(c_4 s) \cos(c_5 s)) \cdot V(\tilde{t}, \tilde{x}, \tilde{y}, \tilde{z}) \\ &\quad + c_1^s c_2^{-s} \cos(c_4 s) \sin(c_5 s) \cdot W(\tilde{t}, \tilde{x}, \tilde{y}, \tilde{z}), \\ \tilde{v} &= c_1^s c_2^{-s} \sin(c_3 s) \cos(c_4 s) \cdot U(\tilde{t}, \tilde{x}, \tilde{y}, \tilde{z}) \\ &\quad + c_1^s c_2^{-s} \cos(c_3 s) \cos(c_4 s) \cdot V(\tilde{t}, \tilde{x}, \tilde{y}, \tilde{z}) \\ &\quad + c_1^s c_2^{-s} \cdot \sin(c_4 s) \cdot W(\tilde{t}, \tilde{x}, \tilde{y}, \tilde{z}), \\ \tilde{w} &= -c_1^s c_2^{-s} (\cos(c_3 s) \sin(c_5 s) + \sin(c_3 s) \sin(c_4 s) \cos(c_5 s)) \cdot U(\tilde{t}, \tilde{x}, \tilde{y}, \tilde{z}) \\ &\quad + c_1^s c_2^{-s} (\sin(c_3 s) \sin(c_5 s) - \cos(c_3 s) \sin(c_4 s) \cos(c_5 s)) \cdot V(\tilde{t}, \tilde{x}, \tilde{y}, \tilde{z}) \\ &\quad + c_1^s c_2^{-s} \cos(c_4 s) \cos(c_5 s) \cdot W(\tilde{t}, \tilde{x}, \tilde{y}, \tilde{z}), \\ \tilde{p} &= c_6^{-s} \cdot P(\tilde{t}, \tilde{x}, \tilde{y}, \tilde{z}), \end{aligned}$$

$$\tilde{q} = c_1^{-2s} c_2^{2s} c_6^{-s} . Q(\tilde{t}, \tilde{x}, \tilde{y}, \tilde{z}),$$

where $\tilde{t} = c_1^s . t$, and

$$\begin{aligned} \tilde{x} = & c_2^s (\sin(c_5 s) \sin(c_4 s) \sin(c_3 s) + \cos(c_5 s) \cos(c_3 s)) . x \\ & + c_2^s (\sin(c_3 s) \cos(c_5 s) - \cos(c_3 s) \sin(c_4 s) \sin(c_5 s)) . y \\ & - c_2^s \cos(c_4 s) \sin(c_5 s) . z, \end{aligned}$$

$$\tilde{y} = -c_2^s \sin(c_3 s) \cos(c_4 s) . x + (c_2 s) \cos(c_3 s) \cos(c_4 s) . y - c_2^s \sin(c_4 s) . z,$$

$$\begin{aligned} \tilde{z} = & c_2^s (\cos(c_3 s) \sin(c_5 s) - \sin(c_3 s) \sin(c_4 s) \cos(c_5 s)) . x \\ & + c_2^s (\cos(c_3 s) \sin(c_4 s) \cos(c_5 s) + \sin(c_3 s) \sin(c_5 s)) . y \\ & + c_2^s \cos(c_4 s) \cos(c_5 s) . z. \end{aligned}$$

The construction of the group-invariant solutions for each of the one-dimensional subgroups in the optimal system proceeds in the same fashion.

6. Acknowledgements

I am grateful to professor Ian M. Anderson for his consideration and his mathematical career to the development the "DifferentialGeometry" package of MAPLE 11. It is an applicable tool to extremely complex and tedious geometrical computations.

References

- [1] G.W. BLUMAN, J.D. COLE, *Similarity Methods for Differential Equations*, Springer, Berlin, 1974.
- [2] N.H. IBRAGIMOV, *Transformation Groups Applied to Mathematical Physics*, Reidel, Dordrecht, 1985.
- [3] M. NADJAFIKHAH, A. MAHDIPOUR-SHIRAYEH, *Galilean Classification of Spacetime Curves*, arXiv e-print: arXiv:0711.1979.
- [4] F. OLIVERI, M.P. SPECIALE, *Exact solutions to the unsteady equations of perfect gases through Lie group analysis and substitution principles*, Int. J. Non-Linear Mech. 37 (2002) 257–274.
- [5] P.J. OLVER, *Applications of Lie Groups to Differential Equations*, Springer, New York, 1986.
- [6] P.J. OLVER, *Equivalence, Invariants, and Symmetry*, Cambridge University Press, 1995.

- [7] L.V. OVSIANNIKOV, *Group Analysis of Differential Equations*, Academic Press, New York, 1982.
- [8] T.R. SEKHAR, V.D. SHARMA, *Similarity solutions for three dimensional Euler equations using Lie group analysis*, Appl. Math. and Comp. 196 (2008) 147–157.
- [9] W.H. STEEB, *Continuous symmetries, Lie algebras, differential equations and computer algebra*, World Sci. Pub. Co., Singapore, 1996.

Group analysis of three dimensional Euler equations of gas dynamics

Mehdi Nadjafikhah*

Abstract

In this paper, the equations governing the unsteady flow of a perfect polytropic gas in three space dimensions are considered. The basic similarity reductions for this system are performed. Reduced equations and exact solutions associated with the symmetries are obtained. This results is used to give the invariance of system up to Galilean motions of space-times $\mathbb{R}^4$. Then, an optimal system of one-dimensional sub-algebras for symmetry algebra of this system is given.

Keywords: Euler equations, Lie group of transformations, Symmetry generators, Similarity solution, optimal system of Lie sub-algebras.

Mathematics Subject Classification: 35C05, 76M60.

1 Introduction

The equations governing the unsteady flow of a perfect polytropic gas in three space dimensions are

$$\begin{aligned}
 \frac{\partial u}{\partial t} + u \frac{\partial u}{\partial x} + v \frac{\partial u}{\partial y} + w \frac{\partial u}{\partial z} + \frac{1}{q} \frac{\partial p}{\partial x} &= 0, \\
 \frac{\partial v}{\partial t} + u \frac{\partial v}{\partial x} + v \frac{\partial v}{\partial y} + w \frac{\partial v}{\partial z} + \frac{1}{q} \frac{\partial p}{\partial y} &= 0, \\
 \frac{\partial w}{\partial t} + u \frac{\partial w}{\partial x} + v \frac{\partial w}{\partial y} + w \frac{\partial w}{\partial z} + \frac{1}{q} \frac{\partial p}{\partial z} &= 0, \\
 \frac{\partial q}{\partial t} + q \left(\frac{\partial u}{\partial x} + \frac{\partial v}{\partial y} + \frac{\partial w}{\partial z} \right) + u \frac{\partial q}{\partial x} + v \frac{\partial q}{\partial y} + w \frac{\partial q}{\partial z} &= 0, \\
 \frac{\partial p}{\partial t} + \gamma p \left(\frac{\partial u}{\partial x} + \frac{\partial v}{\partial y} + \frac{\partial w}{\partial z} \right) + u \frac{\partial p}{\partial x} + v \frac{\partial p}{\partial y} + w \frac{\partial p}{\partial z} &= 0.
 \end{aligned} \tag{1}$$

where t is the time and x, y, z are the space coordinates; $q(t, x, y, z)$ is the density, $p(t, x, y, z)$ the pressure, $u(t, x, y, z)$, $v(t, x, y, z)$ and $w(t, x, y, z)$

*Faculty of Mathematics, Iran University of Science and Technology, Narmak, Tehran, I.R.IRAN. e-mail: m_nadjafikhah@iust.ac.ir

the velocity components in the x , y and z directions, respectively, and $\gamma \in \mathbb{R}$ is the adiabatic index. These equations are called "*three dimensional Euler equations of gas dynamics*" [8] and section 6.3 of [9].

In this paper, we consider the equations governing the unsteady flow of a perfect polytropic gas in three space dimensions. The basic similarity reductions for this system are performed. Reduced equations and exact solutions associated with the symmetries are obtained. We find an optimal system of one-dimensional sub-algebras for symmetry algebra of this system.

In section 2 we find the full symmetry algebra of system (1). Chapter 3 is devoted to the group-invariant solutions to the system. The structure of full symmetry algebra illustrated in section. The last section deals with a optimal system of sub-algebras.

2 Lie symmetry of the system

The classical method for finding symmetry reductions of PDE is the Lie group method of infinitesimal transformations. To apply the classical method to (1), we consider the one-parameter Lie group of infinitesimal transformations in (t, x, y, z, u, v, p, q) given by

$$\begin{aligned}\tilde{t} &= t + s.\xi_1(t, x, y, z, u, v, p, q) + O(s^2), \\ \tilde{x} &= x + s.\xi_2(t, x, y, z, u, v, p, q) + O(s^2), \\ \tilde{y} &= y + s.\xi_3(t, x, y, z, u, v, p, q) + O(s^2), \\ \tilde{z} &= z + s.\xi_4(t, x, y, z, u, v, p, q) + O(s^2), \\ \tilde{u} &= u + s.\eta_1(t, x, y, z, u, v, p, q) + O(s^2), \\ \tilde{v} &= v + s.\eta_2(t, x, y, z, u, v, p, q) + O(s^2), \\ \tilde{w} &= w + s.\eta_3(t, x, y, z, u, v, p, q) + O(s^2), \\ \tilde{p} &= p + s.\eta_4(t, x, y, z, u, v, p, q) + O(s^2), \\ \tilde{q} &= q + s.\eta_5(t, x, y, z, u, v, p, q) + O(s^2).\end{aligned}\tag{2}$$

where s is the group parameter. One requires that this transformation leaves the sub-manifold S invariant. Let

$$X = \xi_1 \partial_t + \xi_2 \partial_x + \xi_3 \partial_y + \xi_4 \partial_z + \eta_1 \partial_u + \eta_2 \partial_v + \eta_3 \partial_w + \eta_4 \partial_p + \eta_5 \partial_q$$

be the corresponding infinitesimal transformation; where coefficients are real-valued C^∞ -functions of $t, x, y, z, u, v, w, p, q$. The infinitesimal criterion of invariance (1) will thus involve $t, x, y, z, u, v, w, p, q$ and the derivatives of u, v, w, p, q with respect to t, x, y, z , as well as $\xi_i, i = 1, 2, 3, 4$, and $\eta_j, j = 1, 2, 3, 4, 5$, and their partial derivatives. After eliminating any dependencies among the derivatives of the m's caused by the system itself, we can then equate the coefficients of the remaining unconstrained partial derivatives of u to zero. This will result in a large number of elementary partial differential equations for the coefficient functions ξ_i, η_j of the infinitesimal generator, called the determining equations for the symmetry

group of the given system. In this case, we find the determining equations for the symmetry group of the system (1) to be the following:

$$\begin{aligned}
\xi_{1,t^2} &= \xi_{1,x} = \xi_{1,y} = \xi_{1,z} = \xi_{1,u} = \xi_{1,v} = \xi_{1,w} = \xi_{1,q} = \xi_{1,p} = 0, \\
\xi_{2,x} &= \xi_{2,u} = \xi_{2,v} = \xi_{2,w} = \xi_{2,q} = \xi_{2,p} = \xi_{2,t^2} = \xi_{2,ty} = \xi_{2,y^2} = 0, \\
\xi_{3,y} &= \xi_{3,u} = \xi_{3,v} = \xi_{3,w} = \xi_{3,q} = \xi_{3,p} = \xi_{3,t^2} = 0, \\
\xi_{4,z} &= \xi_{4,w} = \xi_{4,q} = \xi_{4,p} = 0, \\
\xi_{4,t^2} &= \xi_{4,tx} = \xi_{4,ty} = \xi_{4,x^2} = \xi_{4,xy} = \xi_{4,y^2} = 0, \\
\xi_{3,x} &= \xi_{2,y}, \quad \xi_{3,z} + \xi_{4,y} = 0, \quad \xi_{4,v} = \xi_{4,u}, \quad \xi_{2,z} + \xi_{4,x} = 0,
\end{aligned} \tag{3}$$

and

$$\begin{aligned}
\eta_1 &= -u\xi_{1,t} + \xi_{2,t} + v\xi_{2,y} - w\xi_{4,x}, \\
\eta_2 &= -w\xi_{4,y} + \xi_{3,t} - u\xi_{2,y} - v\xi_{1,t}, \\
\eta_3 &= u\xi_{4,x} + \xi_{4,t} + v\xi_{4,y} - w\xi_{1,t}, \\
\eta_4 &= q\eta_5/p + 2q\xi_{1,t}, \\
\eta_{5,t} &= \eta_{5,x} = \eta_{5,y} = \eta_{5,z} = \eta_{5,u} = \eta_{5,v} = \eta_{5,w} = \eta_{5,q} = 0, \\
\eta_5 &= p\eta_{5,p}.
\end{aligned} \tag{4}$$

First, equations (3) require that ξ_i s are just

$$\begin{aligned}
\xi_1 &= a_1t + a_2, \\
\xi_2 &= a_3t + a_4x + a_5y + a_6z + a_7, \\
\xi_3 &= a_8t - a_5x + a_4y + a_9z + a_{10}, \\
\xi_4 &= a_{11}t - a_6x - a_9y + a_4z + a_{12},
\end{aligned} \tag{5}$$

where $a_1, \dots, a_{12}$ are arbitrary constants. Then, equations (4) and (5) requires that

$$\begin{aligned}
\eta_1 &= (a_4 - a_1)u + a_5v - a_6w + a_3, \\
\eta_2 &= -a_5u + (a_4 - a_1)v + a_4w + a_8, \\
\eta_3 &= -a_6u - a_9v + (a_4 - a_1)w + a_{11}, \\
\eta_4 &= (2a_1 - 2a_4 + a_{13})q, \\
\eta_5 &= a_{13}p,
\end{aligned} \tag{6}$$

where a_{13} is an arbitrary constant. Therefore

Theorem 1. *The Lie algebra of infinitesimal symmetries of the system (1) is spanned by the 13 vector fields*

$$\begin{aligned}
X_1 &= \partial_x, \\
X_2 &= \partial_y, \\
X_3 &= \partial_z, \\
X_4 &= \partial_t, \\
X_5 &= t \partial_x + \partial_u, \\
X_6 &= t \partial_y + \partial_v, \\
X_7 &= t \partial_z + \partial_w, \\
X_8 &= t \partial_t - u \partial_u - v \partial_v - w \partial_w + 2q \partial_q, \\
X_9 &= x \partial_x + y \partial_y + z \partial_z + u \partial_u + v \partial_v + w \partial_w - 2r \partial_r, \\
X_{10} &= y \partial_x - x \partial_y + v \partial_u - u \partial_v, \\
X_{11} &= -z \partial_y + y \partial_z - w \partial_v + v \partial_w, \\
X_{12} &= -z \partial_x + x \partial_z - w \partial_u + u \partial_w, \\
X_{13} &= q \partial_q + p \partial_p,
\end{aligned}
\tag{7}$$

These infinitesimal symmetries will generate a Lie algebra $\mathfrak{g}$ over the field of real or complex numbers. The commutator table of Lie algebra $\mathfrak{g}$ for (1) is given below, where the entry in the i th row and j th column is defined as $[X_i, X_j] = X_i.X_j - X_j.X_i$, $i, j = 1, \dots, 13$.

	X_1	X_2	X_3	X_4	X_5	X_6	X_7	X_8	X_9	X_{10}	X_{11}	X_{12}	X_{13}
X_1	0	0	0	0	0	0	0	0	X_1	$-X_2$	0	X_3	0
X_2	0	0	0	0	0	0	0	0	X_2	X_1	X_3	0	0
X_3	0	0	0	0	0	0	0	0	X_3	0	$-X_2$	$-X_1$	0
X_4	0	0	0	0	X_1	X_2	X_3	X_4	0	0	0	0	0
X_5	0	0	0	$-X_1$	0	0	0	$-X_5$	X_5	$-X_6$	0	X_7	0
X_6	0	0	0	$-X_2$	0	0	0	$-X_6$	X_6	X_5	X_7	0	0
X_7	0	0	0	$-X_3$	0	0	0	$-X_7$	X_7	0	$-X_6$	$-X_5$	0
X_8	0	0	0	$-X_4$	X_5	X_6	X_7	0	0	0	0	0	0
X_9	$-X_1$	$-X_2$	$-X_3$	0	$-X_5$	$-X_6$	$-X_7$	0	0	0	0	0	0
X_{10}	X_2	$-X_1$	0	0	X_6	$-X_5$	0	0	0	0	$-X_{12}$	X_{11}	0
X_{11}	0	$-X_3$	X_2	0	0	$-X_7$	X_6	0	0	X_{12}	0	$-X_{10}$	0
X_{12}	$-X_3$	0	X_1	0	$-X_7$	0	X_5	0	0	$-X_{11}$	X_{10}	0	0
X_{13}	0	0	0	0	0	0	0	0	0	0	0	0	0

3 Reduction of the system

Now, we look at the group-invariant solutions to the system (1).

The system (1) is a sub-manifold M of jet space $J^2(\mathbb{R}^4, \mathbb{R}^5)$ defined by the following equations

$$u_t + uu_x + vu_y + wu_z + p_x/q = 0,$$

$$\begin{aligned}
v_t + uv_x + vv_y + wv_z + p_y/q &= 0, \\
w_t + uw_x + vw_y + ww_z + p_z/q &= 0, \\
q_t + q(u_x + v_y + w_z) + uq_x + vq_y + wq_z &= 0, \\
p_t + \gamma p(u_x + v_y + w_z) + up_x + vp_y + wp_z &= 0.
\end{aligned} \tag{8}$$

Doing as section 3.1 of [6] and find (in a sense) the most general group-invariant solutions to the system (1).

Theorem 2. *The one-parameter groups $g_i(s) : M \rightarrow M$ generated by the X_i , $i = 1, \dots, 13$ are given in the following table:*

$$\begin{aligned}
g_1 &: (t, x + s, y, z, u, v, w, p, q) \\
g_2 &: (t, x, y + s, z, u, v, w, p, q) \\
g_3 &: (t, x, y, z + s, u, v, w, p, q) \\
g_4 &: (t + s, x, y, z, u, v, w, p, q) \\
g_5 &: (t, x + st, y, z, u + s, v, w, p, q) \\
g_6 &: (t, x, y + st, z, u, v + s, w, p, q) \\
g_7 &: (t, x, y, z + st, u, v, w + s, p, q) \\
g_8 &: (e^s t, x, y, z, e^{-s} u, e^{-s} v, e^{-s} w, p, e^{2s} q) \\
g_9 &: (t, e^s x, e^s y, e^s z, e^s u, e^s v, e^s w, p, e^{-2s} q) \\
g_{10} &: (t, x \cos s + y \sin s, y \cos s - x \sin s, z, \\
&\quad v \sin s + u \cos s, v \cos s - u \sin s, w, p, q) \\
g_{11} &: (t, x, y \cos s - z \sin s, z \cos s + y \sin s, \\
&\quad u, v \cos s - w \sin s, w \cos s + v \sin s, p, q) \\
g_{12} &: (t, x \cos s - z \sin s, y, x \sin s + z \cos s, \\
&\quad u \cos s - w \sin s, v, w \cos s + u \sin s, p, q) \\
g_{13} &: (t, x, y, z, u, v, w, e^s p, e^s q)
\end{aligned} \tag{9}$$

Where, entries give the transformed point

$$\exp(sX_i)(t, x, y, z, u, v, w, p, q) = (\tilde{t}, \tilde{x}, \tilde{y}, \tilde{z}, \tilde{u}, \tilde{v}, \tilde{w}, \tilde{p}, \tilde{q}).$$

Since each group g_i , $i = 1, \dots, 13$, is a symmetry group of (1), then

Theorem 3. *Let $i = 1, \dots, 13$, $s \in \mathbb{R}$, and $\lambda \in \mathbb{R} - \{0\}$. If*

$$\begin{aligned}
u &= U(t, x, y, z), \quad v = V(t, x, y, z), \quad w = W(t, x, y, z), \\
p &= P(t, x, y, z), \quad q = Q(t, x, y, z),
\end{aligned}$$

is a solution of the system (1), so are the functions

$$\begin{aligned} u^{(i)} &= U^{(i)}(t, x, y, z), \quad v^{(i)} = V^{(i)}(t, x, y, z), \quad w^{(i)} = W^{(i)}(t, x, y, z), \\ p^{(i)} &= P^{(i)}(t, x, y, z), \quad q^{(i)} = Q^{(i)}(t, x, y, z), \end{aligned}$$

where

$$\begin{aligned} u^{(1)} &= U(t, x + s, y, z), & u^{(2)} &= U(t, x, y + s, z), \\ v^{(1)} &= V(t, x + s, y, z), & v^{(2)} &= V(t, x, y + s, z), \\ w^{(1)} &= W(t, x + s, y, z), & w^{(2)} &= W(t, x, y + s, z), \\ p^{(1)} &= P(t, x + s, y, z), & p^{(2)} &= P(t, x, y + s, z), \\ q^{(1)} &= Q(t, x + s, y, z), & q^{(2)} &= Q(t, x, y + s, z). \end{aligned}$$

$$\begin{aligned} u^{(3)} &= U(t, x, y, z + s), & u^{(4)} &= U(t + s, x, y, z), \\ v^{(3)} &= V(t, x, y, z + s), & v^{(4)} &= V(t + s, x, y, z), \\ w^{(3)} &= W(t, x, y, z + s), & w^{(4)} &= W(t + s, x, y, z), \\ p^{(3)} &= P(t, x, y, z + s), & p^{(4)} &= P(t + s, x, y, z), \\ q^{(3)} &= Q(t, x, y, z + s), & q^{(4)} &= Q(t + s, x, y, z). \end{aligned}$$

$$\begin{aligned} u^{(5)} &= U(t, x + st, y, z) - s, & u^{(6)} &= U(t, x, y + st, z), \\ v^{(5)} &= V(t, x + st, y, z), & v^{(6)} &= V(t, x, y + st, z) - s, \\ w^{(5)} &= W(t, x + st, y, z), & w^{(6)} &= W(t, x, y + st, z), \\ p^{(5)} &= P(t, x + st, y, z), & p^{(6)} &= P(t, x, y + st, z), \\ q^{(5)} &= Q(t, x + st, y, z), & q^{(6)} &= Q(t, x, y + st, z). \end{aligned}$$

$$\begin{aligned} u^{(7)} &= U(t, x, y, z + st), & u^{(8)} &= \lambda.U(\lambda.t, x, y, z), \\ v^{(7)} &= V(t, x, y, z + st), & v^{(8)} &= \lambda.V(\lambda.t, x, y, z), \\ w^{(7)} &= W(t, x, y, z + st) - s, & w^{(8)} &= \lambda.W(\lambda.t, x, y, z), \\ p^{(7)} &= P(t, x, y, z + st), & p^{(8)} &= P(\lambda.t, x, y, z), \\ q^{(7)} &= Q(t, x, y, z + st), & q^{(8)} &= \frac{1}{\lambda^2}.Q(\lambda.t, x, y, z). \end{aligned}$$

$$\begin{aligned} u^{(9)} &= \frac{1}{\lambda}.U(t, \lambda.x, \lambda.y, \lambda.z), & u^{(13)} &= U(t, x, y, z), \\ v^{(9)} &= \frac{1}{\lambda}.V(t, \lambda.x, \lambda.y, \lambda.z), & v^{(13)} &= V(t, x, y, z), \\ w^{(9)} &= \frac{1}{\lambda}.W(t, \lambda.x, \lambda.y, \lambda.z), & w^{(13)} &= W(t, x, y, z), \\ p^{(9)} &= P(t, \lambda.x, \lambda.y, \lambda.z), & p^{(13)} &= \lambda.P(t, x, y, z), \\ q^{(9)} &= \lambda^2.Q(t, \lambda.x, \lambda.y, \lambda.z), & q^{(13)} &= \lambda.Q(t, x, y, z). \end{aligned}$$

$$\begin{aligned} u^{(10)} &= \cos s.U(t, x \cos s + y \sin s, y \cos s - x \sin s, z) \\ &\quad - \sin s.V(t, x \cos s + y \sin s, y \cos s - x \sin s, z), \\ v^{(10)} &= \sin s.U(t, x \cos s + y \sin s, y \cos s - x \sin s, z) \\ &\quad + \cos s.V(t, x \cos s + y \sin s, y \cos s - x \sin s, z), \end{aligned}$$

$$\begin{aligned}
w^{(10)} &= W(t, x \cos s + y \sin s, y \cos s - x \sin s, z), \\
p^{(10)} &= P(t, x \cos s + y \sin s, y \cos s - x \sin s, z), \\
q^{(10)} &= Q(t, x \cos s + y \sin s, y \cos s - x \sin s, z). \\
\\
u^{(11)} &= U(t, x, y \cos s - z \sin s, z \cos s + y \sin s), \\
v^{(11)} &= \cos s.V(t, x, y \cos s - z \sin s, z \cos s + y \sin s) \\
&\quad + \sin s.W(t, x, y \cos s - z \sin s, z \cos s + y \sin s), \\
w^{(11)} &= \cos s.W(t, x, y \cos s - z \sin s, z \cos s + y \sin s) \\
&\quad - \sin s.V(t, x, y \cos s - z \sin s, z \cos s + y \sin s), \\
p^{(11)} &= P(t, x, y \cos s - z \sin s, z \cos s + y \sin s), \\
q^{(11)} &= Q(t, x, y \cos s - z \sin s, z \cos s + y \sin s). \\
\\
u^{(12)} &= \cos s.U(t, x \cos s - z \sin s, y, x \sin s + z \cos s) \\
&\quad + \sin s.W(t, x \cos s - z \sin s, y, x \sin s + z \cos s), \\
v^{(12)} &= V(t, x \cos s - z \sin s, y, x \sin s + z \cos s), \\
w^{(12)} &= \cos s.W(t, x \cos s - z \sin s, y, x \sin s + z \cos s) \\
&\quad - \sin s.U(t, x \cos s - z \sin s, y, x \sin s + z \cos s), \\
p^{(12)} &= P(t, x \cos s - z \sin s, y, x \sin s + z \cos s), \\
q^{(12)} &= Q(t, x \cos s - z \sin s, y, x \sin s + z \cos s).
\end{aligned}$$

4 Structure of Lie algebra $\mathfrak{g}$

In this section, we determine the structure of full symmetry algebra $\mathfrak{g}$. of system (1).

The center $\mathfrak{z}$ of $\mathfrak{g}$ is $\text{Span}_{\mathbb{R}}\{X_{13}\}$. Therefore, the quotient algebra $\mathfrak{g}_1 := \mathfrak{g}/\mathfrak{z}$ is $\text{Span}_{\mathbb{R}}\{Y_1, \dots, Y_{12}\}$; where $Y_i := X_i + \mathfrak{z}$, $i = 1, \dots, 12$. The commutator table of Lie algebra $\mathfrak{g}_1$ is given below, where the entry in the i th row and j th column is defined as $[Y_i, Y_j] = Y_i.Y_j - Y_j.Y_i$,

$i, j = 1, \dots, 12$.

	Y_1	Y_2	Y_3	Y_4	Y_5	Y_6	Y_7	Y_8	Y_9	Y_{10}	Y_{11}	Y_{12}
Y_1	0	0	0	0	0	0	0	0	Y_1	$-Y_2$	0	Y_3
Y_2	0	0	0	0	0	0	0	0	Y_2	Y_1	Y_3	0
Y_3	0	0	0	0	0	0	0	0	Y_3	0	$-Y_2$	$-Y_1$
Y_4	0	0	0	0	Y_1	Y_2	Y_3	Y_4	0	0	0	0
Y_5	0	0	0	$-Y_1$	0	0	0	$-Y_5$	Y_5	$-Y_6$	0	Y_7
Y_6	0	0	0	$-Y_2$	0	0	0	$-Y_6$	Y_6	Y_5	Y_7	0
Y_7	0	0	0	$-Y_3$	0	0	0	$-Y_7$	Y_7	0	$-Y_6$	$-Y_5$
Y_8	0	0	0	$-Y_4$	Y_5	Y_6	Y_7	0	0	0	0	0
Y_9	$-Y_1$	$-Y_2$	$-Y_3$	0	$-Y_5$	$-Y_6$	$-Y_7$	0	0	0	0	0
Y_{10}	Y_2	$-Y_1$	0	0	Y_6	$-Y_5$	0	0	0	0	$-Y_{12}$	Y_{11}
Y_{11}	0	$-Y_3$	Y_2	0	0	$-Y_7$	Y_6	0	0	Y_{12}	0	$-Y_{10}$
Y_{12}	$-Y_3$	0	Y_1	0	$-Y_7$	0	Y_5	0	0	$-Y_{11}$	Y_{10}	0

The algebra $\mathfrak{g}_1$ is not solvable, because

$$\mathfrak{g}_1^{(1)} := [\mathfrak{g}_1, \mathfrak{g}_1] = \text{Span}_{\mathbb{R}}\{Y_1, \dots, Y_{10}\}, \text{ and } \mathfrak{g}_1^{(2)} := [\mathfrak{g}_1^{(1)}, \mathfrak{g}_1^{(1)}] = \mathfrak{g}_1^{(1)}.$$

Lie algebra $\mathfrak{g}_1$ admits a Levi-decomposition into the semi-direct sum

$$\mathfrak{g}_1 = \mathfrak{r} \ltimes \mathfrak{s},$$

where $\mathfrak{r} = \text{Span}_{\mathbb{R}}\{Y_1, \dots, Y_9\}$ is the radical of $\mathfrak{g}_1$ (the largest solvable ideal contained in $\mathfrak{g}_1$) and $\mathfrak{s} = \text{Span}_{\mathbb{R}}\{Y_{10}, Y_{11}, Y_{12}\}$ is a semi-simple subalgebra of $\mathfrak{g}_1$. The radical $\mathfrak{r}$ is uniquely defined but the semi-simple subalgebra $\mathfrak{s}$ is not. $\mathfrak{s}$ as an algebra is isomorphic to 3-dimensional special orthogonal algebra $\mathfrak{so}(3) := \{A \in \text{Mat}(3 \times 3; \mathbb{R}) \mid A^t = -A\}$, which is a simple algebra.

The radical $\mathfrak{r}$ is solvable, with and

$$\mathfrak{r}^{(1)} = \text{Span}_{\mathbb{R}}\{Y_1, \dots, Y_7\}, \quad \mathfrak{r}^{(2)} = \text{Span}_{\mathbb{R}}\{Y_1, Y_2, Y_3\}, \quad \mathfrak{r}^{(3)} = \{0\}.$$

It is semi-direct sum $\mathfrak{r} = \mathfrak{r}^{(1)} \ltimes \mathfrak{g}_2$, where $\mathfrak{g}_2 = \text{Span}_{\mathbb{R}}\{Y_8, Y_9\}$ is isomorphic to the two dimensional Abelian Lie algebra $\mathbb{R}^2$; i.e.

$$\mathfrak{r} = \mathfrak{r}^{(1)} \ltimes \mathbb{R}^2.$$

$\mathfrak{r}^{(1)}$ is semi-direct sum $\mathfrak{r}^{(1)} = \mathfrak{r}^{(2)} \ltimes \mathfrak{g}_3$, where $\mathfrak{g}_3 = \text{Span}_{\mathbb{R}}\{Y_4, \dots, Y_7\}$ is isomorphic to the four dimensional Abelian Lie algebra $\mathbb{R}^4$, and $\mathfrak{r}^{(2)}$ is isomorphic to the three dimensional Abelian Lie algebra $\mathbb{R}^3$; i.e.

$$\mathfrak{r}^{(1)} = \mathbb{R}^3 \ltimes \mathbb{R}^4.$$

Theorem 4. *The symmetry algebra $\mathfrak{g}$ of system (1) have the following structure*

$$\mathfrak{g} \cong \mathbb{R} \times ((\mathbb{R}^3 \ltimes \mathbb{R}^4) \ltimes \mathbb{R}^2) \ltimes \mathfrak{so}(3).$$

Theorem 5. *There is a normal Lie-subalgebra of symmetry Lie-group of system (1) which is isomorphic to the Galilean group of $\mathbb{R}^4$:*

$$\text{Gal}(4, \mathbb{R}) = \left\{ \begin{pmatrix} 1 & \mathbf{0} & s \\ \mathbf{v} & \mathbf{R} & \mathbf{y} \\ 0 & \mathbf{0} & 1 \end{pmatrix} \mid \mathbf{R} \in \text{O}(3, \mathbb{R}), s \in \mathbb{R}, \text{ and } \mathbf{y}, \mathbf{v} \in \mathbb{R}^3 \right\}.$$

Therefore, the system (1) is invariant up to Galilean motions of space-times $\mathbb{R}^4$ (see [3]).

Proof. Let $\mathfrak{b} = \text{Span}_{\mathbb{R}}\{X_1, \dots, X_7, X_{10}, X_{11}, X_{12}\}$. Lie algebra structure of $\mathfrak{b}$ is as $\mathfrak{gal}(3)$, the lie algebra of Galilean group $\text{Gal}(4, \mathbb{R})$, and $\mathfrak{b}$ an ideal of $\mathfrak{g}$. Therefore, there is a Lie-subgroup of G such that its Lie-algebra is $\mathfrak{b}$, by the Theorem 2.53 of Olver [6]. $\square$

Conclusion 1. $\mathfrak{g}$ is semi-direct sum of

$$\mathfrak{gal}(4) \cong \text{Span}_{\mathbb{R}}\{X_1, \dots, X_7, X_{10}, X_{11}, X_{12}\}$$

and 3-dimensional Abelian algebra $\mathbb{R}^3 \cong \text{Span}\{X_8, X_9, X_{13}\}$.

5 Optimal system of sub-algebras

As is well known, the Lie group theoretic method plays an important role in finding exact solutions and performing symmetry reductions of differential equations. Since any linear combination of infinitesimal generators is also an infinitesimal generator, there are always infinitely many different symmetry subgroups for the differential equation. So, a mean of determining which subgroups would give essentially different types of solutions is necessary and significant for a complete understanding of the invariant solutions. As any transformation in the full symmetry group maps a solution to another solution, it is sufficient to find invariant solutions which are not related by transformations in the full symmetry group, this has led to the concept of an optimal system [7]. The problem of finding an optimal system of subgroups is equivalent to that of finding an optimal system of subalgebras. For one-dimensional subalgebras, this classification problem is essentially the same as the problem of classifying the orbits of the adjoint representation. This problem is attacked by the naive approach of taking a general element in the Lie algebra and subjecting it to various adjoint transformations so as to simplify it as much as possible. The idea of using the adjoint representation to classify group-invariant solutions was due to [7] and [5].

The adjoint action is given by the Lie series

$$\text{Ad}(\exp(sX_i)X_j) = X_j - s[X_i, X_j] + \frac{s^2}{2}[X_i, [X_i, X_j]] - \dots, \quad (10)$$

where $[X_i, X_j]$ is the commutator for the Lie algebra, s is a parameter, and $i, j = 1, \dots, 13$. We can write the adjoint action for the Lie algebra $\mathfrak{g}$, and show that

Theorem 6. *A one-dimensional optimal system of (1) is given by*

- 1) $X = X_8 + \sum_{i=9}^{13} a_i X_i,$
- 2) $X = a_7 X_7 + X_8 + X_9 + \sum_{i=10}^{13} a_i X_i,$
- 3) $X = a_6 X_6 + a_7 X_7 + X_8 + X_9 + \sum_{i=11}^{13} a_i X_i,$
- 4) $X = a_5 X_5 + X_8 + X_9 + a_{11} X_i + a_{13} X_{13},$
- 5) $X = \sum_{i=5}^7 a_i X_i + X_8 + X_9 + a_{13} X_{13},$
- 6) $X = a_4 X_4 + X_9 + \sum_{i=10}^{13} a_i X_i,$
- 7) $X = X_4 + a_5 X_5 + a_8 X_8 + \sum_{i=10}^{13} a_i X_i,$
- 8) $X = X_4 + a_6 X_6 + a_8 X_8 + a_{10} X_{10} + a_{12} X_{12} + a_{13} X_{13},$
- 9) $X = X_4 + a_7 X_7 + a_8 X_8 + a_{10} X_{10} + a_{12} X_{12} + a_{13} X_{13},$
- 10) $X = X_4 + \sum_{i=5}^8 a_i X_i + a_{13} X_{13},$
- 11) $X = X_4 + a_7 X_7 + \sum_{i=10}^{13} a_i X_i,$
- 12) $X = X_7 + \sum_{i=10}^{13} a_i X_i,$
- 13) $X = a_3 X_3 + \sum_{i=10}^{13} a_i X_i,$
- 14) $X = X_4 + a_6 X_6 + \sum_{i=11}^{13} a_i X_i,$
- 15) $X = X_4 + a_5 X_5 + a_{11} X_{11} + a_{13} X_{13},$
- 16) $X = X_4 + \sum_{i=5}^7 a_i X_i + a_{13} X_{13},$
- 17) $X = X_6 + \sum_{i=11}^{13} a_i X_i,$
- 18) $X = a_2 X_2 + \sum_{i=11}^{13} a_i X_i,$
- 19) $X = a_2 X_2 + a_{12} X_{12} + a_{13} X_{13},$
- 20) $X = X_5 + a_{11} X_{11} + a_{13} X_{13},$
- 21) $X = a_2 X_2 + a_3 X_3 + X_5 + a_{13} X_{13},$
- 22) $X = a_1 X_1 + X_{11} + a_{13} X_{13},$
- 23) $X = a_1 X_1 + a_3 X_3 + X_6 + a_{13} X_{13},$
- 24) $X = a_1 X_1 + a_2 X_2 + a_3 X_3 + a_{13} X_{13},$
- 25) $X = X_5 + a_{10} X_{10} + a_{11} X_{11} + a_{13} X_{13},$
- 26) $X = a_3 X_3 + X_{10} + a_{11} X_{11} + a_{13} X_{13}.$

Proof: $F(sX_i) : \mathfrak{g} \rightarrow \mathfrak{g}$ defined by $X \mapsto \text{Ad}(\exp(sX_i)X)$ is a linear map, for $i = 1, \dots, 13$. The matrix M_{sX_i} of $F(sX_i)$, $i = 1, \dots, 13$, with respect to basis $\{X_1, \dots, X_{13}\}$ is

$$\begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ -s & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & s & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & -s & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix},$$

[illegible]

[illegible]

[illegible]

[illegible]

[illegible]

[illegible]

[illegible]

[illegible]

[illegible]

$$\begin{bmatrix} \cos s & 0 & \sin s & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ -\sin s & 0 & \cos s & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & \cos s & 0 & \sin s & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & -\sin s & 0 & \cos s & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & \cos s & \sin s & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & -\sin s & \cos s & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix},$$

and I_{13} , respectively. Let $X = \sum_{i=1}^{13} a_i X_i$, then

$$\begin{aligned} F(s_7.X_7) \circ F(s_6.X_6) \circ \cdots \circ F(s_1.X_1) : X \mapsto \\ (s_3a_{12} - s_2a_{10} - s_1a_9 + a_1 - s_5s_4a_8 + s_5a_4 - s_4a_5).X_1 \\ + (-s_6s_4a_8 + s_3a_{11} - s_2a_9 + s_1a_{10} + a_2 + s_6a_4 - s_4a_6).X_2 \\ + (-s_7s_4a_8 - s_3a_9 - s_2a_{11} - s_1a_{12} + a_3 + s_7a_4 - s_4a_7).X_3 \quad (11) \\ + (a_4 - s_4a_8).X_4 + (-s_6a_{10} - s_5a_9 + s_5a_8 + a_5 + s_7a_{12}).X_5 \\ + (-s_6a_9 + s_6a_8 + s_5a_{10} + a_6 + s_7a_{11}).X_6 \\ + (s_7a_8 - s_6a_{11} - s_5a_{12} + a_7 - s_7a_9).X_7 \\ + a_8.X_8 + a_9.X_9 + a_{10}.X_{10} + a_{11}.X_{11} + a_{12}.X_{12} + a_{13}.X_{13} \end{aligned}$$

If a_8 , and $a_9 \neq 0$, and $a_8 \neq 2a_9$, then we can make the coefficients of $X_1, \dots, X_7$ vanish. Scaling X if necessary, we can assume that $a_8 = 1$. And X is reduced to Case 1.

If $a_8 = a_9 \neq 0$, and $a_{10} \neq 0$, then we can make the coefficients of $X_1, \dots, X_6$ vanish. Scaling X if necessary, we can assume that $a_8 = 1$. And X is reduced to Case 2.

If $a_8 = a_9 \neq 0$, $a_{10} = 0$, and $a_{12} \neq 0$, then we can make the coefficients of $X_1, \dots, X_5$, and X_{10} vanish. Scaling X if necessary, we can assume that $a_8 = 1$. And X is reduced to Case 3.

If $a_8 = a_9 \neq 0$, $a_{10} = a_{12} = 0$, and $a_{11} \neq 0$, then we can make the coefficients of $X_1, \dots, X_4$, X_6 , X_7 , X_{10} , and X_{12} vanish. Scaling X if necessary, we can assume that $a_8 = 1$. And X is reduced to Case 4.

If $a_8 = a_9 \neq 0$, $a_{10} = a_{11} = a_{12} = 0$, then we can make the coefficients of $X_1, \dots, X_4$, and $X_{10}, \dots, X_{11}$ vanish. Scaling X if necessary, we can assume that $a_8 = 1$. And X is reduced to Case 5.

If $a_8 = 0$, and $a_9 \neq 0$, then we can make the coefficients of $X_1, \dots, X_3$, and $X_5, \dots, X_8$ vanish. Scaling X if necessary, we can assume that $a_9 = 1$. And X is reduced to Case 6.

If $a_8 \neq 0$, $a_9 = 0$, and $a_{11} \neq 0$, then we can make the coefficients of $X_1, \dots, X_3$, and X_6 , X_7 , and X_9 vanish. And X is reduced to Case 7.

If $a_8 \neq 0$, $a_9 = a_{11} = 0$, and $a_{12} \neq 0$, then we can make the coefficients of $X_1, \dots, X_3$, and X_5 , X_7 , X_9 , and X_{11} vanish. And X is reduced to Case 8.

If $a_8 \neq 0$, $a_9 = a_{11} = a_{12} = 0$, and $a_{10} \neq 0$, then we can make the coefficients of $X_1, \dots, X_3$, and X_5 , X_6 , X_9 , and X_{11} vanish. And X is reduced to Case 9.

If $a_8 \neq 0$, and $a_9 = a_{10} = a_{11} = a_{12} = 0$, then we can make the coefficients of $X_1, \dots, X_3$, and $X_9, \dots, X_{12}$ vanish. And X is reduced to Case 10.

If $a_8 = a_9 = 0$, $a_4 \neq 0$, and $a_{10} \neq 0$, then we can make the coefficients of $X_1, \dots, X_3$, X_5 , X_6 , X_8 , and X_9 vanish. Scaling X if necessary, we can assume that $a_4 = 1$. And X is reduced to Case 11.

If $a_4 = a_8 = a_9 = 0$, $a_{10} \neq 0$, $a_{12} \neq 0$, and $a_6 a_{12} + a_7 a_{10} \neq a_5 a_{11}$, then we can make the coefficients of $X_1, \dots, X_6$, X_8 , and X_9 vanish. Scaling X if necessary, we can assume that $a_7 = 1$. And X is reduced to Case 12.

If $a_4 = a_8 = a_9 = 0$, $a_{10} \neq 0$, $a_{12} \neq 0$, and $a_6 a_{12} + a_7 a_{10} = a_5 a_{11}$, then we can make the coefficients of X_1 , X_2 , and $X_4, \dots, X_9$ vanish. And X is reduced to Case 13.

If $a_8 = a_9 = a_{10} = 0$, and $a_4 \neq 0$, then we can make the coefficients of $X_1, \dots, X_3$, X_5 , and $X_7, \dots, X_{10}$ vanish. Scaling X if necessary, we can assume that $a_4 = 1$. And X is reduced to Case 14.

If $a_8 = \dots = a_{10} = 0$, $a_{12} = 0$, $a_4 \neq 0$, and $a_{11} \neq 0$, then we can make the coefficients of $X_1, \dots, X_3$, and $X_8, \dots, X_{10}$ vanish. Scaling X if necessary, we can assume that $a_4 = 1$. And X is reduced to Case 15.

If $a_8 = \dots = a_{12} = 0$, and $a_4 \neq 0$, then we can make the coefficients of $X_1, \dots, X_3$, and $X_8, \dots, X_{10}$ vanish. Scaling X if necessary, we can assume that $a_4 = 1$. And X is reduced to Case 16.

If $a_4 = a_8 = a_9 = a_{10} = 0$, $a_{12} \neq 0$, and $a_5 a_{11} \neq a_6 a_{12}$, then we can make the coefficients of $X_1, \dots, X_5$, and $X_7, \dots, X_{10}$ vanish. Scaling X if necessary, we can assume that $a_6 = 1$. And X is reduced to Case 17.

If $a_4 = a_8 = a_9 = a_{10} = 0$, $a_{12} \neq 0$, $a_5 a_{11} = a_6 a_{12}$, and $a_6 \neq 0$, then we can make the coefficients of $X_1, \dots, X_5$, and $X_7, \dots, X_{10}$ vanish. And X is reduced to Case 18.

If $a_4 = a_6 = a_8 = a_9 = a_{10} = 0$, $a_{12} \neq 0$, $a_5 a_{11} = a_6 a_{12}$, and $a_6 \neq 0$, then we can make the coefficients of $X_1, \dots, X_5$, and $X_7, \dots, X_{10}$ vanish. And X is reduced to Case 18.

If $a_4 = a_6 = a_8 = \dots = a_{11} = 0$, $a_{12} \neq 0$, $a_5 a_{11} = a_6 a_{12}$, $a_6 \neq 0$, and $a_5 \neq 0$, then we can make the coefficients of X_1 , and $X_3, \dots, X_{11}$ vanish. And X is reduced to Case 19.

If $a_4 = a_8 = a_9 = a_{10} = a_{12} = 0$, $a_5 \neq 0$, and $a_{11} \neq 0$, then we can make the coefficients of $X_1, \dots, a_4$, $X_6, \dots, X_{10}$, and a_{12} vanish. Scaling X if necessary, we can assume that $a_5 = 1$. And X is reduced to Case 20.

If $a_4 = a_8 = \dots = a_{12} = 0$, and $a_5 \neq 0$, then we can make the coefficients of X_1 , a_4 , and $X_8, \dots, X_{12}$ vanish. Scaling X if necessary, we can assume that $a_5 = 1$. And X is reduced to Case 21.

If $a_4 = a_5 = a_8 = a_9 = a_{10} = a_{12} = 0$, and $a_{11} \neq 0$, then we can make the coefficients of $X_2, \dots, X_{10}$ and a_{12} vanish. Scaling X if necessary, we can assume that $a_{11} = 1$. And X is reduced to Case 22.

If $a_4 = a_5 = a_8 = \dots = a_{12} = 0$, and $a_6 \neq 0$, then we can make the coefficients of X_2 , a_5 , and $X_7, \dots, X_{12}$ vanish. Scaling X if necessary, we can assume that $a_6 = 1$. And X is reduced to Case 23.

If $a_4 = a_5 = a_6 = 0$, and $a_8 = \dots = a_{12} = 0$, then we can make the coefficients of $X_4, \dots, X_{12}$ vanish. And X is reduced to Case 24.

If $a_4 = a_8 = a_9 = a_{12} = 0$, $a_{10} \neq 0$, and $a_7 a_{10} \neq a_5 a_{11}$, then we can make the coefficients of $X_1, \dots, X_4$, $X_6, \dots, X_9$, and $a_{12} = 0$ vanish. Scaling X if necessary, we can assume that $a_5 = 1$. And X is reduced to Case 25.

If $a_4 = a_8 = a_9 = a_{12} = 0$, $a_7 a_{10} = a_5 a_{11}$, and $a_{10} \neq 0$, then we can make the coefficients of $X_1, \dots, X_4$, $X_6, \dots, X_9$, and $a_{12} = 0$ vanish. Scaling X if necessary, we can assume that $a_{10} = 1$. And X is reduced to Case 26. And the Theorem follows. $\square$

According to our optimal system of one-dimensional subalgebras of the full symmetry algebra $\mathfrak{g}$, we need only find group-invariant solutions for 26 one-parameter subgroups generated by X as Theorem 6. For example, as a direct consequence of Theorem 3 and case 1 of Theorem 6, we deduce the following conclusion:

Conclusion 2 *Let $s, c_1, \dots, c_6 \in \mathbb{R}$ with c_1, c_2 , and $c_6 \neq 0$, and s be sufficiently small. Then, if*

$$\begin{aligned} u &= U(t, x, y, z), \quad v = V(t, x, y, z), \quad w = W(t, x, y, z), \\ p &= P(t, x, y, z), \quad q = Q(t, x, y, z), \end{aligned}$$

is a solution of the system (1), so are the functions

$$\begin{aligned} \tilde{u} &= c_1^s c_2^{-s} (\cos(c_3 s) \cos(c_5 s) - \sin(c_3 s) \sin(c_4 s) \sin(c_5 s)) \cdot U(\tilde{t}, \tilde{x}, \tilde{y}, \tilde{z}) \\ &\quad - c_1^s c_2^{-s} (\sin(c_3 s) \cos(c_5 s) + \cos(c_3 s) \sin(c_4 s) \cos(c_5 s)) \cdot V(\tilde{t}, \tilde{x}, \tilde{y}, \tilde{z}) \\ &\quad + c_1^s c_2^{-s} \cos(c_4 s) \sin(c_5 s) \cdot W(\tilde{t}, \tilde{x}, \tilde{y}, \tilde{z}), \\ \tilde{v} &= c_1^s c_2^{-s} \sin(c_3 s) \cos(c_4 s) \cdot U(\tilde{t}, \tilde{x}, \tilde{y}, \tilde{z}) \\ &\quad + c_1^s c_2^{-s} \cos(c_3 s) \cos(c_4 s) \cdot V(\tilde{t}, \tilde{x}, \tilde{y}, \tilde{z}) \\ &\quad + c_1^s c_2^{-s} \cdot \sin(c_4 s) \cdot W(\tilde{t}, \tilde{x}, \tilde{y}, \tilde{z}), \\ \tilde{w} &= -c_1^s c_2^{-s} (\cos(c_3 s) \sin(c_5 s) + \sin(c_3 s) \sin(c_4 s) \cos(c_5 s)) \cdot U(\tilde{t}, \tilde{x}, \tilde{y}, \tilde{z}) \\ &\quad + c_1^s c_2^{-s} (\sin(c_3 s) \sin(c_5 s) - \cos(c_3 s) \sin(c_4 s) \cos(c_5 s)) \cdot V(\tilde{t}, \tilde{x}, \tilde{y}, \tilde{z}) \\ &\quad + c_1^s c_2^{-s} \cos(c_4 s) \cos(c_5 s) \cdot W(\tilde{t}, \tilde{x}, \tilde{y}, \tilde{z}), \\ \tilde{p} &= c_6^{-s} \cdot P(\tilde{t}, \tilde{x}, \tilde{y}, \tilde{z}), \\ \tilde{q} &= c_1^{-2s} c_2^{2s} c_6^{-s} \cdot Q(\tilde{t}, \tilde{x}, \tilde{y}, \tilde{z}), \end{aligned}$$

where $\tilde{t} = c_1^s t$, and

$$\begin{aligned} \tilde{x} &= c_2^s (\sin(c_5 s) \sin(c_4 s) \sin(c_3 s) + \cos(c_5 s) \cos(c_3 s)) \cdot x \\ &\quad + c_2^s (\sin(c_3 s) \cos(c_5 s) - \cos(c_3 s) \sin(c_4 s) \sin(c_5 s)) \cdot y \\ &\quad - c_2^s \cos(c_4 s) \sin(c_5 s) \cdot z, \\ \tilde{y} &= -c_2^s \sin(c_3 s) \cos(c_4 s) \cdot x + (c_2 s) \cos(c_3 s) \cos(c_4 s) \cdot y - c_2^s \sin(c_4 s) \cdot z, \\ \tilde{z} &= c_2^s (\cos(c_3 s) \sin(c_5 s) - \sin(c_3 s) \sin(c_4 s) \cos(c_5 s)) \cdot x \end{aligned}$$

$$\begin{aligned}
&+c_2^s(\cos(c_3s)\sin(c_4s)\cos(c_5s)+\sin(c_3s)\sin(c_5s)).y \\
&+c_2^s\cos(c_4s)\cos(c_5s).z.
\end{aligned}$$

The construction of the group-invariant solutions for each of the one-dimensional subgroups in the optimal system proceeds in the same fashion.

6 Acknowledgements

I am grateful to professor Ian M. Anderson for his consideration and his mathematical career to the development the "DifferentialGeometry" package of MAPLE 11. It is an applicable tool to extremely complex and tedious geometrical computations.

References

- [1] G.W. BLUMAN, J.D. COLE, *Similarity Methods for Differential Equations*, Springer, Berlin, 1974.
- [2] N.H. IBRAGIMOV, *Transformation Groups Applied to Mathematical Physics*, Reidel, Dordrecht, 1985.
- [3] M. NADJAFIKHAH, A. MAHDIPOUR-SHIRAYEH, *Galilean Classification of Spacetime Curves*, arXiv e-print: arXiv:0711.1979.
- [4] F. OLIVERI, M.P. SPECIALE, *Exact solutions to the unsteady equations of perfect gases through Lie group analysis and substitution principles*, Int. J. Non-Linear Mech. 37 (2002) 257–274.
- [5] P.J. OLVER, *Applications of Lie Groups to Differential Equations*, Springer, New York, 1986.
- [6] P.J. OLVER, *Equivalence, Invariants, and Symmetry*, Cambridge University Press, 1995.
- [7] L.V. OVSIANNIKOV, *Group Analysis of Differential Equations*, Academic Press, New York, 1982.
- [8] T.R. SEKHAR, V.D. SHARMA, *Similarity solutions for three dimensional Euler equations using Lie group analysis*, Appl. Math. and Comp. 196 (2008) 147–157.
- [9] W.H. STEEB, *Continuous symmetries, Lie algebras, differential equations and computer algebra*, World Sci. Pub. Co., Singapore, 1996.

This sentence no verb.

Check for spaces (there should be none between ‘for’ and ‘spaces’), and around inclusion of proper texts (one space between ‘around’ and ‘inclusion’, and one between ‘inclusion’ and ‘of’).

Als check for multiple paragraphs.

Such as here. (‘Such as’ should be a new paragraph.)

This text is rather small don’t you think? (The ‘rather small’ should be footnotesize.)

And now we disable environment (in between ‘disable’ and ‘environment’ is some excluded material).

We switch to **bold face inside** a comment. (The phrase ‘bold face inside’ should be bold.)

This is a line of text

****Comment: 1 line(s) removed****another line of text

****Comment: 3 line(s) removed****last line of text.

$$\begin{aligned}\omega_1 &:= dt \\ \omega_2 &:= \frac{dx}{f} \\ \omega_3 &:= dy - \frac{g}{f}d(x)\end{aligned}$$

$$\begin{aligned}d\omega_1 &= -f_x\omega_1\wedge\omega_3 - f_y\omega_2\wedge\omega_3, \\ d\omega_2 &= -g_x\omega_1\wedge\omega_3 - g_y\omega_2\wedge\omega_3, \\ d\omega_3 &= 0.\end{aligned}$$

$$\Phi : \bar{t} = \phi(t), \quad \bar{x} = \psi(x,y,t), \quad \bar{y} = \eta(x,y,t).$$

$$\Phi^*\left(\begin{array}{c}\bar{\omega}_1 \\ \bar{\omega}_2 \\ \bar{\omega}_3\end{array}\right)=\left(\begin{array}{ccc}a_1&0&0 \\ a_2&a_3&0 \\ 0&0&a_4\end{array}\right)\left(\begin{array}{c}\omega_1 \\ \omega_2 \\ \omega_3\end{array}\right)$$

$$G=\left\{g=\left(\begin{array}{ccc}a_1&0&0 \\ a_2&a_3&0 \\ 0&0&a_4\end{array}\right):a_i\in\mathbb{R},\,a_1a_3a_4\neq0\right\}$$

$$\left(\begin{array}{c}\bar{\theta}_1 \\ \bar{\theta}_2 \\ \bar{\theta}_3\end{array}\right)=\left(\begin{array}{ccc}a_1&0&0 \\ a_2&a_3&0 \\ 0&0&a_4\end{array}\right)\left(\begin{array}{c}\omega_1 \\ \omega_2 \\ \omega_3\end{array}\right)=\left(\begin{array}{c}a_1\omega_1 \\ a_2\omega_1+a_3\omega_2 \\ a_4\omega_3\end{array}\right)$$

$$dg\cdot g^{-1}=\left(\begin{array}{ccc}\frac{da_1}{a_1}&0&0 \\ \frac{da_2}{a_1}-\frac{a_2\,da_3}{a_1a_3}&\frac{da_3}{a_3}&0 \\ 0&0&\frac{da_4}{a_4}\end{array}\right)$$

$$\pi_1:=\frac{da_1}{a_1},\;\pi_2:=\frac{da_2}{a_1}-\frac{a_2\,da_3}{a_1a_3},\;\pi_3:=\frac{da_3}{a_3},\;\pi_4:=\frac{da_4}{a_4}.$$

$$\begin{aligned}
d\theta_1 &= \pi_1 \wedge \theta_1 + \frac{a_2 f_y - a_3 f_x}{a_3 a_4} \theta_1 \wedge \theta_3 - \frac{a_1 f_y}{a_3 a_4} \theta_2 \wedge \theta_3 \\
d\theta_2 &= \pi_2 \wedge \theta_1 + \pi_3 \wedge \theta_2 + \frac{-a_2 a_3 f_x + a_2^2 f_y - a_3^2 g_x + a_2 a_3 g_y}{a_1 a_3 a_4} \theta_1 \wedge \theta_3 - \frac{a_2 f_y + a_3 g_y}{a_3 a_4} \theta_2 \wedge \theta_3 \\
d\theta_3 &= \pi_4 \wedge \theta_3
\end{aligned}$$

The coefficient $T = -\frac{a_1 f_y}{a_3 a_4}$ in $d\theta_1$ is an essential torsion. By assuming $T = -1$, we find that $a_1 = a_3 a_4 / f_y$.

$$\begin{aligned}
\omega_1 &:= dx - f(x, y) dt \\
\omega_2 &:= dy - g(x, y) dt \\
\omega_3 &:= dt
\end{aligned}$$

$$\begin{aligned}
d\omega_1 &= -f_x \omega_1 \wedge \omega_3 - f_y \omega_2 \wedge \omega_3, \\
d\omega_2 &= -g_x \omega_1 \wedge \omega_3 - g_y \omega_2 \wedge \omega_3, \\
d\omega_3 &= 0.
\end{aligned}$$

$$\Phi : \bar{t} = \phi(t), \quad \bar{x} = \psi(x, y, t), \quad \bar{y} = \eta(x, y, t).$$

$$\Phi^* \begin{pmatrix} \bar{\omega}_1 \\ \bar{\omega}_2 \\ \bar{\omega}_3 \end{pmatrix} = \begin{pmatrix} a_1 & 0 & 0 \\ a_2 & a_3 & 0 \\ 0 & 0 & a_4 \end{pmatrix} \begin{pmatrix} \omega_1 \\ \omega_2 \\ \omega_3 \end{pmatrix}$$

$$G = \left\{ g = \begin{pmatrix} a_1 & 0 & 0 \\ a_2 & a_3 & 0 \\ 0 & 0 & a_4 \end{pmatrix} : a_i \in \mathbb{R}, a_1 a_3 a_4 \neq 0 \right\}$$

$$\begin{pmatrix} \bar{\theta}_1 \\ \bar{\theta}_2 \\ \bar{\theta}_3 \end{pmatrix} = \begin{pmatrix} a_1 & 0 & 0 \\ a_2 & a_3 & 0 \\ 0 & 0 & a_4 \end{pmatrix} \begin{pmatrix} \omega_1 \\ \omega_2 \\ \omega_3 \end{pmatrix} = \begin{pmatrix} a_1 \omega_1 \\ a_2 \omega_1 + a_3 \omega_2 \\ a_4 \omega_3 \end{pmatrix}$$

$$dg \cdot g^{-1} = \begin{pmatrix} \frac{da_1}{a_1} & 0 & 0 \\ \frac{da_2}{a_1} - \frac{a_2 da_3}{a_1 a_3} & \frac{da_3}{a_3} & 0 \\ 0 & 0 & \frac{da_4}{a_4} \end{pmatrix}$$

$$\pi_1 := \frac{da_1}{a_1}, \quad \pi_2 := \frac{da_2}{a_1} - \frac{a_2 da_3}{a_1 a_3}, \quad \pi_3 := \frac{da_3}{a_3}, \quad \pi_4 := \frac{da_4}{a_4}.$$

$$\begin{aligned}
d\theta_1 &= \pi_1 \wedge \theta_1 + \frac{a_2 f_y - a_3 f_x}{a_3 a_4} \theta_1 \wedge \theta_3 - \frac{a_1 f_y}{a_3 a_4} \theta_2 \wedge \theta_3 \\
d\theta_2 &= \pi_2 \wedge \theta_1 + \pi_3 \wedge \theta_2 + \frac{-a_2 a_3 f_x + a_2^2 f_y - a_3^2 g_x + a_2 a_3 g_y}{a_1 a_3 a_4} \theta_1 \wedge \theta_3 - \frac{a_2 f_y + a_3 g_y}{a_3 a_4} \theta_2 \wedge \theta_3 \\
d\theta_3 &= \pi_4 \wedge \theta_3
\end{aligned}$$

The coefficient $T = -\frac{a_1 f_y}{a_3 a_4}$ in $d\theta_1$ is an essential torsion. By assuming $T = -1$, we find that $a_1 = a_3 a_4 / f_y$.